

(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11) Publication number:

**0 491 077 A1**

(12)

## EUROPEAN PATENT APPLICATION

(21) Application number: **90124775.9**

(51) Int. Cl.<sup>5</sup>: **A61K 39/12, A61K 39/29,  
C12N 15/33, C12N 15/36**

(22) Date of filing: **19.12.90**

(43) Date of publication of application:  
**24.06.92 Bulletin 92/26**

(84) Designated Contracting States:  
**DE**

(71) Applicant: **GA INVESTMENT S.A.**  
**Banco Union Building 6th Floor, Samuel**  
**Lewis Avenue**  
**Panama(PA)**

(72) Inventor: **Thoma, Hans A., Dr.**  
**Giselastrasse 3**  
**W-8000 München 40(DE)**

(74) Representative: **Strehl, Schübel-Hopf,**  
**Groening**  
**Maximilianstrasse 54 Postfach 22 14 55**  
**W-8000 München 22(DE)**

(54) **A composition used as a therapeutic agent against chronic viral hepatic diseases.**

(57) A combination, comprising at least one polypeptide sequence, mediating the antigenicity of one or more epitopes, and a carrier, capable of presenting this/these polypeptide sequence(s), which are useful for the production of a medicament for the treatment of chronic viral hepatitis, is provided.

EP 0 491 077 A1

The present invention relates to a composition comprising a polypeptide sequence prepared by recombinant DNA techniques and a carrier to provide a curing agent against chronic viral hepatic diseases. The invention relates to DNA sequences coding for said polypeptide sequences and to transfected cells for the expression of the same.

At least five different viruses, namely Hepatitis virus A, B, C, D and E, are able to trigger the clinical aspect of an acute hepatitis. Hepatitis A and E, which are transferred enterically, always heal, whereas hepatitis B, C (formerly called parenteral hepatitis Non-A Non-B), and D can progress into a chronic stage of inflammation, which in turn can result in liver cirrhosis and primary hepatocellular carcinoma.

There is relatively little data available on hepatitis C and D, on methods for the diagnosis and their treatment and on the respective viruses. The hepatitis D virus is a RNA virus which is known to be incomplete. Therefore, it needs a helper virus to develop in patients and is found only in individuals infected with HBV. Only very recently the hepatitis C virus has been detected, and an antibody test (anti-HCV) facilitating the diagnosis of chronic hepatitis C infections has been developed. However, there is an increasingly urgent need for a treatment to cure this disease.

The same holds true for chronic hepatitis B, a much better studied disease with respect to its recognition by immunological methods, its causative virus and the viral life cycle and DNA sequence. Patients are said to be chronic carriers of the hepatitis B virus if the viral DNA persists longer than ten weeks, the HBe-antigen (HBeAg) for more than 12 weeks, or if the hepatitis B surface-antigen (HBsAg) is persistent longer than six months.

Roughly three hundred million people are deemed to suffer from chronic hepatitis B, most of them living in the Far East.

For these people the main risk to be infected appears to be during or immediately after birth, since a chronically infected mother transfers the virus to her newborn. 90 percent of the children infected this way will become chronically infected, too, during later life. In the Western World infection occurs more commonly later in life, during childhood or even adulthood, mainly by a parenteral or sexual transmission. In these cases of hepatitis B infection after birth only five to ten percent of the infected become chronic carriers. The virus transferred, however, is not responsible for distinct reactions shown by infected people either to eliminate the virus or to retain it in the body lifelong. Consequently, it seems to be a matter of the immunological status that determines the future physical condition.

The HB-virion (Dane particle) is composed of different structural proteins, the core proteins and the surface (S) proteins. The latter are translation products of an open reading frame encompassing the coding sequence of three S-type domains, each of which starts with an ATG triplet capable of initiating translation in vivo. The domains are referred to as preS1, preS2 and S in the order of 5' to the 3'end of the molecule. There are six protein products derived from this ORF: a glycosylated and a non-glycosylated form of the major protein (gp27 and p24) translated from the S domain only (226 amino acids), a middle protein (281 amino acids) having one or two polysaccharide side chains (gp33 and gp36, respectively), that is encoded by the preS2- and S-region, and finally, both a glycosylated (gp42) and a non-glycosylated (p39) form of the large protein (389-400 amino acids, depending upon the viral serotype), which is formed by translation of preS1, preS2 and S. The core proteins are HBcAg and HBeAg, the latter one conceivably being a processing product of HBcAg.

The Dane particle, which is the infectious virion, comprises both core and surface proteins, whereas the filaments consist of a mixture of the six surface antigens. The S peptides alone assemble to form the so-called 20nm particles, which are completely uninfected.

Patients infected by the HB virus pass through several stages of the hepatitis, before they are regarded to be chronically HBV-infected. Immediately after infection an infectious stage will follow, characterized by the presence of HBeAg in the serum. Continued HBs antigenaemia in spite of inhibited HBV replication indicates the presence of viral DNA sequences integrated into the cellular genome of the patient. The integrated viral sequences do not enable the host cell to synthesize the complete virus. However, liver cells having HBV-sequences integrated are capable of producing HBsAg only, which in turn is detectable in the serum of the patient and is an indicator for chronic hepatitis B. Most probably the transformed hepatocytes are not lysed by cytotoxic T-cells, but proliferate and induce either chronic persistent hepatitis (CPH) or chronic active hepatitis (CAH), which may then proceed to cirrhosis of the liver or to primary hepatocellular carcinoma resulting in premature death of the patient.

Recently it has been established that patients who are chronically HBV infected show a defect in endogenous interferon production (Abb et al., 1985: J. Med. Virol 16. 171-176). This was the rationale to treat patients suffering from chronic hepatitis B, as indicated by the presence of HBeAg and HBV-DNA in the serum, with interferon  $\alpha$  (IFN $\alpha$ ). Controlled trials with large numbers of patients showed that the administration of interferon  $\alpha$  resulted in significantly increased elimination of the hepatitis B-virus, when

compared to controls. However, persons infected at or around time of birth do not appear to seroconvert in response to this therapy. This phenomenon unfortunately precludes some 75% of carriers from IFN $\alpha$  therapy.

At present, the exact mode of action of interferon  $\alpha$  on chronic hepatitis B remains unclear. Its antiviral activity might protect infected cells from infection or reduce viral transcription, translation and replication in HBV-infected cells. Interferon further has immunomodulatory effects by activating T-cells, macrophages and NK-cells and by inducing the expression of MHC class I proteins.

Another approach to treat chronic hepatitis B is based on the idea to inhibit replication of the virus, thus impairing its defence sufficiently to render the host immune system capable of eliminating the virus. This led to test antiviral drugs such as adenine arabinoside and adenine arabinoside monophosphate for treatment of chronically HBV infected individuals. However, less than half of the patients responded to this therapy, either by sustained or transient seroconversion (HBeAg<sup>+</sup> to anti-HBe<sup>+</sup>). A further negative aspect of these antiviral drugs are their immunosuppressive properties. Other drugs that have been tested for treatment of chronic carriers include interferon  $\beta$  and , acycloguanosine (acyclovir), interleukin 2, steroids, such as prednisolone, and combinations thereof. But none of them could provide better results than treatment with interferon  $\alpha$ . Only a combination therapy, including the initial administration of steroids followed by that of IFN  $\alpha$  may increase the response rate in selected patients.

It is known from the prior art, that chronically HBV infected chimpanzees can not be cured by treatment with HBsAg (bound to a tetanus toxoid) nor with anti-HBs antibodies. Furthermore, it has been attempted to immunize chronically HBV infected patients by administration of S peptides. This treatment did not even result in anti-HBs antibody formation in these persons.

Additionally, according to the definition, chronic carriers of hepatitis B virus are characterized in that HBsAg is detectable in their serum. Therefore, it has been absolutely unforeseeable, that a combination, comprising a T-cell activating epitope of the viral S peptide, according to the present invention, is able to induce an immunization in and a final healing of chronic carriers of hepatitis virus B.

Considering the above-discussed state of the art it is the objective of the present invention to provide an effective therapeutic agent for the treatment of viral chronic hepatic diseases which leads to a complete response (i.e. to the sustained inhibition of HBV-replication, the loss of HBV DNA and DNA polymerase and to a decrease and finally the disappearance of HBeAg and HBsAg in the serum of patients).

According to the present invention this goal is achieved by a combination of a) at least one polypeptide sequence mediating the antigenicity of one or more epitopes and b) a carrier, capable of presenting the epitope sequence(s) a), wherein the polypeptide sequence(s) a) can be bound to carrier b) by adsorption, any chemical bonding or secondary valences.

This invention is furthermore directed to the use of this combination for the production of a medicament for the treatment of chronic viral hepatitis.

It is important that polypeptide sequence a), which may be one or more different polypeptides, mediates the antigenicity of a T cell-activating epitope in a direct or indirect way. According to the present invention polypeptide sequence(s) a) may be a polypeptide or a combination of two or more polypeptides of hepatitis B virus of any subtype, particularly adw, ayw, adr and ady.

These peptides derived from hepatitis B virus may be HBV peptides preS1, preS2 or S or the HBV core antigens.

Useful as polypeptide sequence(s) a) are furthermore any of the above-stated polypeptides or a combination of two or more polypeptides which are modified either by amino acid deletions, whereby at least one epitope comprising at least six consecutive amino acid residues must be preserved, or by adding further amino acids either at the N-terminus, the C-terminus or as insertions into the polypeptide sequence(s) a). In each of these cases it is essential, however, that the biological activity is maintained.

Preferably, polypeptide sequence(s) a) is myristylated.

In order to display the appropriate pharmacological activity it is necessary that in the combination of the present invention polypeptide sequence(s) a) is presented on a carrier b). This carrier consists of a particular substance which for example may consist of particles of a hydrophobic polymer, of inorganic particles, or of particles of a polysaccharide. Preferably, carrier b) is a second polypeptide sequence which forms particles upon secretion, said particles having preferably a diameter of at least 10nm.

It is preferred that the particle forming polypeptide sequence b) is a substantial part of or the complete amino acid sequence of a polypeptide which may be selected from HBV S peptide, HBV core antigen, HAV core antigen, HAV surface antigen, HIV surface antigen and HIV core antigen as well as the surface antigen of polio virus. Preferred as the particle-forming carrier b) is HBV S peptide and/or core peptide.

When used as the carrier sequence b) the above-stated polypeptides may be modified by arbitrary deletions of amino acids, by substitutions of one or more amino acids or by adding one or more amino

acids either at the N-terminus, the C-terminus or by insertion of one or more amino acids into the polypeptide sequence b), provided that the particle-forming capacity is maintained. Preferably, polypeptide sequence b) is myristylated.

If the carrier b) is a polypeptide sequence, both sequences a) and b) may be linked via one or more of the following interactions: hydrophobic anchoring (mediated by myristic acid), disulfide bridge formation, or both sequences may be connected by a peptide bond to form a fusion peptide. In the latter case optionally a spacer sequence may be inserted between polypeptide sequence(s) a) and polypeptide sequence b), which spacer sequence is linked to both polypeptides via peptide bonds.

The present invention furthermore provides a recombinant DNA molecule coding for a combination, that is useful for production of a medicament to treat chronic viral hepatic diseases. The recombinant DNA molecule comprises at least one first DNA sequence, optionally a second, a third and/or a fourth DNA sequence wherein

- i) said at least one first DNA sequence codes for at least one polypeptide sequence a) as defined above,
- ii) said second DNA sequence codes for a polypeptide sequence b) according to the above definition of the particle forming peptide,
- iii) said third DNA sequence codes for a spacer sequence, and
- iv) said fourth DNA sequence codes for a selection marker,

and wherein the DNA sequences are controlled by DNA elements essential for expression, and optionally have a common reading frame.

On account of the fact, that many amino acids are designated by more than one triplet, there exist several DNA sequences embraced by the present invention, which code for the above-defined peptide sequences a) and b). Apart from this, the invention further embraces recombinant DNA molecules, which differ from the above-defined recombinant DNA molecules by the fact, that up to 30% of the nucleotides may be substituted.

A further object of the present application is to provide a host cell transfected with a recombinant DNA molecule coding for the above combination, which is useful for treatment of chronically HBV-infected patients. This host cell may be a mammalian, a yeast or a bacterial cell. For the purpose of this invention it is preferred, that this cell does not produce any human serum proteins or any primate serum proteins other than the polypeptide(s) being comprised within the above combination.

The term "HBV S peptide" as used herein refers to the peptide encoded by the entire region of the HBV genome. The term "HBV pre-S2 peptide" as used herein refers to the peptide encoded by the entire pre-S2 and S regions of the HBV genome. The term "HBV pre-S1 peptide" as used herein refers to the polypeptide encoded by the entire pre-S1, pre-S2 and S regions of the HBV genome. The term "epitope" as used herein refers to a sequence of at least six consecutive amino acids encoded by the designated genome region (e.g. a "HBV pre-S2 epitope" refers to a sequence of at least six amino acids encoded by the pre-S2 region of the HBV genome). The term "T-cell epitope" as used herein refers to an epitope that interacts with receptors on the surface of T-cells to enhance or otherwise effect an immune response.

As used herein "antigenicity" means the ability to provoke an immune response (e.g. acting as an antigen), the ability to cause the production of antibodies (e.g. acting as an antigen) and/or the ability to interact with a cell surface receptor so as to enhance an immune response or production of antibodies.

The term "HBV" means any subtype of the virus, particularly adw, ayw, adr and ayr, described in the literature (P. Valenzuela, Nature Vol. 280, p. 815 (1979), Gerlich, EP-A-85 111 361, Neurath, EP-A-85 102 250). Examples of peptide sequences thereof, constituting polypeptide sequence(s) a), which mediate the antigenicity of one or more epitopes, are shown in the Sequence Listing (SEQ ID No. 17-20, 22).

Preferred embodiments of the present invention are the following combinations:

- HB S-antigen particles with specific epitopes (determinants) of the pre-S1-, pre-S2-, and/or core peptides;
- HB core-antigen particles with specific epitopes (determinants) of the pre-S1-, pre-S2-, S-peptide, and/or of the core antigens;
- Hepatitis A-antigen particles with specific epitopes (determinants) of the hepatitis B S-, pre-S1-, pre-S2-, and/or core-peptides.

Recombinant DNA molecules preferred for the present invention are characterized by the presence of sequences coding for polypeptide sequence(s) a), mediating the antigenicity of one or more T-cell epitopes, and for polypeptide b), which upon secretion forms particles having a diameter of 10nm or more, both of which are under control of a suitable promoter. As examples for sequences coding for a) there may be mentioned any of the sequences listed under ID numbers 1 to 24 in the sequence Listing. Examples for the DNA sequence coding for polypeptide sequence b) are represented by any of the ID-sequences 25 to 27 in the Sequence Listing.

Any of the 24 sequences (ID numbers 1 to 24) may be combined to any sequence disclosed under ID number 25 to 27 in the Sequence Listing, therein both orders a-b and b-a are included.

Hepatitis virus sequences used in the recombinant DNA construct of the present invention can be formed or isolated by any means including isolation and ligation of restriction fragments, chemical synthesis of oligonucleotides using a synthesizer (Cyclon, Bio-Search), and synthesis by the PCR method (T.J. White, N. Arnheim, H. E. Erlich, 1989; The Polymerase Chain Reaction, Technical Focus 5 (6)).

Preferred recombinant DNA molecules were formed by the ligation of synthetic oligonucleotides to a 5' XbaI-BglII 3' fragment (ID number 27) from the S region of the HBV genome, which is derived from a BglII-BglII HBV fragment including the entire pre-S1-pre-S2-S-region, or to the entire S-region. Oligonucleotides used in making such constructs are summarized in Table I below.

Table I

Function	Definition	SEQ ID No.
core (adw)	aa* 59-87	6
core (adw)	aa 2-28	7
core (adw)	aa -10-28	8
core (adw)	aa 29-58	9
core (adw)	aa 1-87	10
core (adw)	aa -10-87	11
core (adw)	aa 70-110	12
core (adw)	aa 80-125	13
core (adw)	aa 88-120	15
S1 (ayw)	aa 9-28	17
S1 (ayw)	aa 83-103	18
S1 (ayw)	aa 20-40	19
S1 (ayw)	aa 59-94	20
S1 (adw)	aa 94-114	21
S1 (adw)	aa 70-105	22
S2 (ayw)	aa 2-21	23
S2 (ayw)	aa 14-33	24

\* aa = amino acid

Other preferred DNA molecules were formed by ligation of core sequences, which are prepared by the PCR method and which code for T-cell epitopes, to the core sequence of HBV (SEQ ID NO 25) functioning as polypeptide sequence b). Oligonucleotides used in preparing these constructs are given in Table II-1.

Table II-1

Function	Definition	SEQ ID No.
core	complete, bp 1901-2500	1
core	C-terminal deletion, bp 1901-2405	2
core	C-terminal deletion and stop codon inserted, bp 1901-2405	3
core/precore	10 aa precore, C-terminal deletion, bp 1871-2405	4
core/precore	10 aa precore, C-terminal deletion and stop codon inserted, bp 1871-2405	5
core	aa (-10-120)	16
core/precore	10 aa precore, complete core, bp 1871-2500	35

Table II-2 shows several examples, where the T-cell epitope-coding DNA sequences have been isolated by restriction fragmentation of the HBV genome and have been ligated to the DNA sequence coding for polypeptide sequence b) as defined above.

Table II-2

Function	Definition	SEQ ID No.
core/precore	complete, bp 1403-31	**
S2 ay/ad		**
S2 (K) ay/ad	S2-S, 7 codons deleted, start codon ATG changed to ATA	14

\*\* Sequence has been published by Galibert, F. et al. (1979: Nature 281, 646-650) and by Ono, Y. et al. (1983: Nucl. Acid Res. 11(6), 1747-1757)

In Table II-3 specific recombinant DNA molecules are listed. The procedure for their construction will be described in more detail in the Examples.

Table II-3

Final construct	T-cell epitope	Particle Former	Selection Gene
MT-core(-10-120) + SAg + neo	core(aa-10-120)	S adw/ayw or S/XbaI/BglII	neo
MT-S1(aa 9-28)-S + egpt	S1(aa 9-28)ay	S adw/ayw or S/XbaI/BglII	egpt#
MT-core-neo	core/precore bp 1403-31	core adw	neo
MT-core(1-87) + HBsAg - neo	core(aa 1-87)	S adw/ayw or S/XbaI/BglII	neo

# egpt = E coli xanthine quanine phosphoribosyl transferase

Preferred recombinant DNA molecules according to the present invention comprise, apart from the regions coding for polypeptides a) and b), an additional DNA sequence coding for a selection marker. Furthermore, they comprise all usual elements essential for the expression, such as promoter sequence, start codon and a polyadenylation signal.

Examples of suitable promoters are the methallothionein (MT), the U2 and the H2K promoter in case of using mammalian cells as a host cell. If yeast or bacterial cells are to be employed, appropriate yeast and bacterial promoters, such as the GCN4- and the GAL 1/10 promoter or the prokaryotic trp- and tac promoters, respectively, may be used.

In order to produce the combination of polypeptide(s) a) and polypeptide b) according to this application the recombinant DNA molecule is inserted into host cells by transfection (in case of mammalian cells), by transformation (in case of yeast and bacterial cells), or by other means. As a host cells of any organism may be used that are capable of transcribing and translating recombinant DNA molecules, such as mammalian, bacterial and yeast cells.

Suitable mammalian cells according to this invention are for example VERO cells (a monkey kidney cell line), 3T3-, C127 and L cells (murine fibroblast cell lines), and CHO (Chinese hamster ovary) cells, which are either positive or negative in dehydrofolate reductase.

According to a specific embodiment of the present invention it is furthermore possible that the above-defined first DNA sequence and the above-defined second DNA sequence, which code for polypeptide sequence(s) a) and for a polypeptide sequence b), respectively, are present in different recombinant DNA molecules, in which case the host cells are cotransfected with both of these recombinant DNA molecules.

Tab. III Possible alternatives of compositions for particles containing T-cell epitopes for targeting chronic hepatitis carrier

	FINAL CONSTRUCT	PROMOTER	SYN	T-CELL-EPIOTOPE		PARTICULAR FORMER	SELECTION GENE	PURIFICATION
				PCR	GENE			
12		MTA202	Core (AA 10 - 87)			entire S adh/gyw S/XbaI/RgII	neo/egpt	
13		MTA202	Core (AA 70 - 110)			entire S adh/gyw S/XbaI/RgII	neo/egpt	
14		MTA202	Core (AA 80 - 125)			entire S adh/gyw S/XbaI/RgII	neo/egpt	
15		MTA202	Core (AA 88 - 120)			entire S adh/gyw S/XbaI/RgII	neo/egpt	
16		MTA202	Core (AA 10 + SAIG + AA 2 - 87)			entire S adh/gyw S/XbaI/RgII	neo/egpt	
17	MT-core (-10 - +120) + SAIG + neo	MTA202		Core (AA 10 - +120)		entire S adh/gyw S/XbaI/RgII	neo/egpt	
18	MT - S1 (AA 9-28) - S + egpt	MTA202	S1 - (AA 9 - 28) (ay)			entire S adh/gyw S/XbaI/RgII	neo/egpt	
19		MTA202	S1 - (AA 83 - 103) (ay)			entire S adh/gyw S/XbaI/RgII	neo/egpt	
20		MTA202	S1 - (AA 20 - 40) (ay)			entire S adh/gyw S/XbaI/RgII	neo/egpt	Materials and Methods
21		MTA202	S1 - (AA 59 - 94) (ay)			entire S adh/gyw S/XbaI/RgII	neo/egpt	
22		MTA202	S1 - (AA 94 - 114) (ad)			entire S adh/gyw S/XbaI/RgII	neo/egpt	
23		MTA202	S1 - (AA 70 - 105) (ad)			entire S adh/gyw S/XbaI/RgII	neo/egpt	
24		MTA202	S1 - (AA 9 - 28) (ay)			core adh	neo/egpt	
25		MTA202	S2 - (AA 2 - 21) (ay)			entire S adh/gyw S/XbaI/RgII	neo/egpt	
26		MTA202	S2 - (AA 14 - 33) (ay)			entire S adh/gyw S/XbaI/RgII	neo/egpt	
27		MTA202			S2 ayw	S ayw/adw	neo/egpt	
28		MTA202	Adapter		S2 - (K) - ayw	S ayw/adw	neo/egpt	

Tab. III Possible alternatives of compositions for particles containing T-cell epitopes for targeting chronic hepatitis carrier

1	FINAL CONSTRUCT	2 PROMOTER	SYN	T-CELL-EPTOPE		GENE	PARTICLE FORMER	SELECTION GENE	3 PURIFICATION
				PCR					
1		MTA202		Core without pre-core			Core (cdw)	neo/epfl	
				e.g. bp 1901 - 2500					
2		MTA202		Core without pre-core, with deletion of the C-terminus			Core (cdw)	neo/epfl	
				e.g. bp 1901 - 2405					
3		MTA202		Core without pre-core, with deletion at the C-terminus + stop signal			Core (cdw)	neo/epfl	
				e.g. bp 1901 - 2405					
4		MTA202		Core and pre-core 10 AA		Core with pre-core i.e. bp 1403 - 31	Core (cdw)	neo/epfl	Materials and Methods
				Core and pre-core 10 AA; with deletion at the C-terminus					
5		MTA202		Core and pre-core 10 AA; with deletion at the C-terminus			Core (cdw)	neo/epfl	
				e.g. bp 1871 - 2405					
6		MTA202		Core and pre-core with deletion at the C-terminus + stop signal			Core (cdw)	neo/epfl	
				e.g. bp 1871 - 2405					
7		MTA202		Core (AA 59 - 87)			entire S adw/aw S/Kba/Bgll	neo/epfl	
8		MTA202		Core (AA 2 - 28)			entire S adw/aw S/Kba/Bgll	neo/epfl	
9		MTA202		Core (AA 10 - 428)			entire S adw/aw S/Kba/Bgll	neo/epfl	
10		MTA202		Core (AA 29 - 58)			entire S adw/aw S/Kba/Bgll	neo/epfl	
11		MTA202		Core (AA 1 - 87) + HBSAg-neo			entire S adw/aw S/Kba/Bgll	neo/epfl	

Notes: 1 see example 3  
2 any of the stated promoters is suitable  
3 see examples

Table III gives an overview on how to combine suitable DNA sequences to get DNA constructs according to the present invention. It is to be noted that any constituents disclosed in this table may be combined to provide a DNA sequence which may be taken, if transfected into a host cell, to produce a combination (comprising polypeptides(s) a) and b)) as a medicament for the treatment of chronic viral hepatitis. The DNA sequences coding for the T-cell epitope sequences have been prepared synthetically (SYN) with a Biosearch Cyclon synthesizer, by PCR procedure (PCR), or by restriction enzyme fragmentation of the viral genome (GENE).

For the treatment of patients suffering from chronic viral hepatitis the combination of polypeptide



sequence(s) a) and a carrier b) may be formulated in any type of a pharmaceutical composition, which furthermore comprises a suitable diluent or pharmaceutical carrier material, such as a buffer solution.

The administration may be effected by any method, i.e. by parenteral (e.g. intravenous or intramuscular) or oral (e.g. by using typhoid bacterial cells to encapsulate the active substance) administration.

5 The pharmaceutical preparation comprises the above-described combination in sufficient concentration to elicit a response upon administration.

#### Brief Description of the Figures

- 10 Figure I shows a DNA construct, coding for a promoter, a particle former sequence and a selection gene (described in Example 3/4).
- Figure II shows a DNA-gene construct containing a promoter, an epitope with the entire HB-S-Ag and a selection gene (described in Example 3/18).
- 15 Figure III shows a DNA construct presenting a promoter, a T-cell epitope with a particle former residue and a selection gene (described in Example 3/21).
- Figure IV shows the AST values of chimpanzee 1 during the Hepa-Care treatment (described in Example 10/1).
- Figure V shows the antigen values of chimpanzee 1 during the Hepa-Care treatment (described in Example 10/1).
- 20 Figure VI shows values of liver enzymes ALT and GGT of chimpanzee 1 booster treated three times with Hepa-Care (described in Example 10/2).
- Figure VII shows values of liver enzymes ALT, AST, and GGT and of antigen of chimpanzee 2 during the Hepa-Care treatment (described in Example 10/3).
- Figure VIII shows the liver enzymes as determined for an untreated control chimpanzee (described in Example 10/3).
- 25 Figures IX & X show the antigen and antibody titers of patient 1 during the Hepa-Care treatment, respectively (described in Example 11).
- Figures XI & XII show the antigen and antibody titers of patient 2 during the Hepa-Care treatment, respectively (described in Example 11).
- 30 Figures XIII & XIV show the antigen and antibody titers of patient 2 during the Hepa-Care treatment, respectively (described in Example 11).
- The invention is more specifically described by the following examples.

#### Example 1

##### 1. Fractionated precipitation with polyethylene glycol (PEG)

35 The supernatant of HBV protein producing cultures was collected and split into portions of 2,400 ml. To each portion 144 g of PEG 6000 (Serva) were added and dissolved by stirring at room temperature for 20 minutes and was stirred for another 6 hours at 4° C. The precipitate was separated by centrifugation in 500 ml bottles in a GS 3 rotor at 9,000 rpm (15,000 x g) for 30 minutes at 10° C. The supernatant was collected and 144 g of PEG 6000 were added and dissolved as described above. The solution was stirred at 4° C for 3 hours. The precipitate from this solution was harvested as described above except that centrifugation was continued for 60 minutes.

##### 2. Gel Chromatography

45 The material obtained after PEG precipitation was redissolved in 20 ml PBS and submitted to gel chromatography on A-5m (BioRad). Column dimensions were 25 x 1000 mm and 480 ml bed volume. In a typical fractionation run 1,000 µg of PEG precipitated HBV protein in 10 to 15 ml was loaded and eluted with PBS at a speed of 6 drops/min (18 ml/h). 3 ml fractions were collected. HBV protein eluted with the first peak. Collected fractions were submitted to a CsCl gradient.

##### 3. Sedimentation in CsCl gradient

55 About 30 fractions covering the first peak in column chromatography on A-5m and containing prepurified HBV protein were collected to approximately 100 ml. This solution was adjusted to a density of 1.30 g/cc with CsCl and subsequently transferred to a polyallomer tube fitting into a SW 27/28 rotor

(Beckman). A gradient was bet by underlaying 4 ml of a CsCl solution of 1.35 g/cc and by overlaying 4 ml of 1.25 g/cc followed by 4 ml of 1.20 g/cc density. This gradient had been run at 28,000 rpm for 50 hours at 10° C. Thereafter the gradient was fractionated, and purified HBV protein floating in the 1.20 g/cc density layer was collected. The solution was desalted by three cycles of dialysis in bags against water.

## Example 2

### Quantitative Determination of HBV protein

#### 1. with Radioimmunoassay

In the AUSRIA II-125 "sandwich" radioimmunoassay (commercially available from Abbot), beads coated with guinea pig antibody to Hepatitis B surface antigen (Anti-HBs) were incubated with serum or plasma or purified protein and appropriate controls. Any HBsAg present was bound to the solid phase antibody. After aspiration of the unbound material and washing of the bead, human 125T-Anti-HBs was allowed to react with the antibody-antigen complex on the bead. The beads were then washed to remove unbound <sup>125</sup>-Anti-HBs.

)-Anti-HBs HBsAg

)-Anti-HBs . HBsAg <sup>125</sup>I-Anti-HBs

)-Anti-HBs . HBsAg . <sup>125</sup>I-Anti-HBs

The radioactivity remaining on the beads was counted in a gamma scintillation counter.

#### 2. with ELISA

In the Enzygnost HBsAg micro "sandwich" assay (commercially available from Behring), wells were coated with anti-HBs. Serum plasma or purified protein and appropriate controls were added to the wells and incubated. After washing, peroxidase-labelled antibodies to HBsAg were reacted with the remaining antigenic determinants. The unbound enzyme-linked antibodies are removed by washing and the enzyme activity on the solid phase was determined. The enzymatically catalyzed reaction of hydrogen peroxide and chromogen was stopped by adding diluted sulfuric acid. The colour intensity was proportional to the HBsAg concentration of the sample and was obtained by photometric comparison of the colour intensity of the unknown samples with the colour intensities of the accompanying negative and positive control sera.

## Example 3

Preparation of gene constructs of the present invention containing promoter, desired antigen sequences and selection gene.

#### 1. Isolation of the MT-promoter.

The plasmid pBPV-342-12 (ATCC 37224) was digested with the endonucleases BglII and BamHI. Three DNA molecules were generated. The fragment of interest contains the methallothionein promoter and a pBR322 sequence comprising 4.5 kb and is easily detectable from the other fragments (2.0 kb and 7.6 kb).

The reaction was performed in a total volume of 200 µl of reaction buffer at a final concentration of 0.5 µg/µl DNA including 100 units of each restriction enzyme. The completion of the digestion was checked after incubation at 37° C for three hours by agarose gel electrophoresis at a 0.8% agarose gel. The reaction was stopped by adding 4 µl 0.5 M EDTA.

The 4.5 kb fragment was separated from the other fragments by preparative 1.2% agarose gel electrophoresis. The DNA was eluted from the agarose gel on DE-81 Whatman filter paper from which the DNA was removed in a high salt buffer. The DNA was purified by a phenol-chloroform extraction and two ethanol precipitations.

#### 2. Ligation of a 1.8 kb fragment coding for the HBV-core-antigen.

A 1.8kb BamHI-BamHI fragment, containing the HBV-core coding regions was isolated from HBV-containing DNA. This fragment was ligated together with the 4.5 kb fragment containing the MT-promoter and the pBR residue (described in 1).

2 µl of the 1.8 kb fragment were mixed with 3 µl of the 4.5 kb fragment and ligated together in a total volume of 10 µl ligation buffer, containing 2 units T4-DNA ligase and 2 mM ATP at 14° C overnight.

The ligation mixture was added to 150 µl competent bacterial cell suspension for DNA up-take. After the DNA up-take the bacterial cells were spread on LB agar plates containing 50 µl/ml ampicillin at volumes of 50 to 300 µl cell suspension per plate. The agar plates were incubated at 37° C overnight. Single isolated bacterial colonies were screened for the presence of a plasmid containing the desired

fragments.

### 3. Screening for desired plasmid containing bacterial colonies.

Single colonies were picked with a toothpick and transferred to a LB-ampicillin medium containing tube (5 ml). The tubes were incubated overnight at 37° in a rapidly shaking environment. A mini-plasmid preparation of each grown bacterial suspension was made. The different resulting DNAs were proved by digestion with the restriction endonuclease BglII. Two molecules were expected, a 400 bp fragment and a 5.9 kb fragment. The digestion was analysed by agarose gel electrophoresis. Plasmid DNA was isolated from the bacterial cells.

### 4. Insertion of a neomycin selection marker.

The plasmid resulting from (3) above was linearized by digestion with the restriction enzyme EcoRI. The reaction was performed in a total volume of 50 µl and a final concentration of 1 µg/µl plasmid DNA. 50 units of EcoRI were added and the digestion was proved after incubation at 37° C for three hours by agarose gel electrophoresis. The reaction was stopped by adding 1 µl of 0.5 M EDTA and the DNA was precipitated with a final concentration of 0.3 M sodium acetate and 3-4 volumes of ethanol at -80° C for 30 minutes. The precipitated DNA was dissolved in 50 µl distilled water.

2 µl of the linearized plasmid was mixed with 3 µl of the DNA fragment containing the methallothionein promoter and the neomycin selection gene (isolated from the plasmid pMT-neo-E (available from ATCC/Exogene) by digestion with the endonuclease EcoRI as a 3.9 kb fragment), and ligated together. Single bacterial colonies were screened for the presence of the desired plasmid.

### 5. Isolation of a fragment containing the U2 promoter sequence.

The plasmid pUC-8-42 (available from Exogene) was digested with the restriction endonucleases EcoRI and Apal. Two DNA molecules were generated. The fragment of interest contains the U2-promoter comprising 340 bp and is easily detectable from the other fragment (3160 bp). The digestion was performed in a total volume of 200 µl reaction buffer at a final concentration of 0.5 µg/µl DNA including 100 units of each restriction enzyme. The completion of the digest was checked after incubation at 37° C for three hours by agarose gel electrophoresis in a 0.7% agarose gel. The reaction was stopped by adding 4 µl 0.5 M EDTA. The 340 bp fragment was separated from the plasmid DNA by preparative 1.2% agarose gel electrophoresis. The DNA was eluted from the agarose gel on DE-81 Whatman filter paper from which the DNA was removed in a high salt buffer. The DNA was purified by a phenol/chloroform extraction and two ethanol precipitations.

### 6. Insertion of the fragment containing the promoter sequence into a polylinker plasmid.

The plasmid pSP165 (commercially available from Promega Biotec) containing a polylinker sequence (containing the following restriction sites: EcoRI, SacI, SmaI, Aval, BamHI, BglII, Sall, PstI, HindIII) was linearized with the restriction enzyme EcoRI.

The reaction was performed in a total volume of 50 µl and a final concentration of 1 µg/µl plasmid DNA. 50 units of EcoRI was added and the digestion was proved after incubation at 37° C for three hours by agarose gel electrophoresis. The reaction was stopped by adding 1 µl of 0.5 M EDTA and the DNA was precipitated with a final concentration of 0.3 M sodium acetate and 3-4 volumes of ethanol at -80° C for 30 minutes. The precipitated DNA was dissolved in 50 µl distilled water.

2 µl of plasmid DNA was mixed with 10 µl of the fragment DNA containing the U2 promoter sequence, and ligated together in a total volume of 25 µl of ligation buffer containing 2 units T4-DNA ligase and 2mM ATP at 14° C overnight. Thereafter, the DNA purified by phenol/chloroform extractions followed by two ethanol precipitations and dissolved in 10 µl distilled water. The resulting sticky ends of EcoRI and Apal had to be converted into blunt ends and ligated. The sticky ends were converted into blunt ends by reaction with the Mung bean nuclease as follows: to 25 µl DNA (1µg/µl concentration) in reaction buffer 20 units of enzyme were added to give a final concentration of 1% glycerol and a final reaction volume of 35 µl. After an incubation for 30 minutes at 30° C the DNA was purified by phenol-chloroform extractions followed by two ethanol precipitations. The DNA was dissolved again in 5 µl of distilled water. The resulting blunt ends were ligated together in 15 µl reaction volume containing 10 x more T4 ligase than used above and 2 mM ATP at 14° C overnight.

The ligation mixture was added to 150 µl competent bacterial cell suspension for DNA up-take. After the DNA up-take the bacterial cells were spread on LB agar plates containing 50 µg/ml ampicillin at volumes of 50 to 300 µl cell suspension per plate. The agar plates were incubated at 37° C overnight. Single isolated bacterial colonies were screened for the presence of a plasmid containing the desired U2-promoter fragment. The resulting plasmid was isolated from the bacterial cells and characterized by restriction enzyme analysis.

### 7. Ligation of synthetic oligo-DNA-nucleotide 89 (SEQ ID No.: 30) together with MT-promoter fragment (4.5 kb).

The 4.5 kb fragment (described in 1) containing the MT-promoter and a pBR residue were ligated together with the synthetic oligonucleotide 89 (SEQ ID No.:30). The ligation mixture was added to 150  $\mu$ l competent bacterial cell suspension for DNA up-take. Single isolated bacterial colonies were screened for the presence of the desired plasmid. The new plasmid was proved by a digestion with the restriction endonucleases EcoRI and XbaI. Two molecules were expected, one 2.0 kb and one 2.6 kb.

8. Ligation of the synthetic oligonucleotide 101 (SEQ ID No.:32) together with plasmid (described in 7).

The plasmid (described in 7) was digested with BglII and BamHI and a fragment of 13 nucleotides was removed (described in 1). The resulting fragment containing the first oligonucleotide 89 (SEQ ID No.:30), was ligated together with oligonucleotide 101 (SEQ ID No.:32), a BglII-BamHI fragment. After DNA up-take single cells were screened for the presence of the desired plasmid. The new plasmid was proved by a digestion with the endonucleases EcoRI and XbaI, or EcoRI and BglII.

9. Ligation of synthetic DNA-oligonucleotide 99 (SEQ ID No.:31) to the 4.5 kb fragment (described in 1).

The 4.5 kb fragment (BglII-BamHI) was ligated together with the DNA oligonucleotide 99 (SEQ ID No.: 31). After screening of single bacterial colonies, containing different DNAs, the desired plasmid was characterized by digestion with EcoRI, resulting in two fragments, 1.9 kb and 2.7 kb, and by positive linearization with BglII or BamHI.

The new plasmid was then digested with PstI and BamHI. Two molecules were expected, one 2.6 kb fragment, containing a pBR residue, the MT-promoter and the oligonucleotide and a 2.0 kb pBR residue. The 2.6 kb fragment was isolated.

10. Ligation of the 2.6 kb fragment of the plasmid described in 9, with a fragment isolated from plasmid (described in 8).

The plasmid (described in 8) containing the DNA oligonucleotides 89 and 101 (SEQ ID No.:30 and 32, respectively) was digested with PstI and BglII. Two fragments were expected. A 2.5kb fragment containing a pBR residue and the MT-promoter and 2.2 kb fragment, containing a pBR residue and both oligos.

This 2.2 kb fragment was ligated together with the 2.6 kb fragment, containing the pBR residue, the MT-promoter and oligo 99 (SEQ ID No.:31) described in 8.

After screening for the desired plasmid, it was characterized by restriction endonuclease digestion with BglII-XbaI. Two fragments were expected, a 270 bp fragment of the oligo-DNA-nucleotides and a 4.5 kb fragment of the MT-promoter and the pBR.

11. Ligation of the 2.3 kb HBV BglII-BglII fragment.

A 2.3 kb BglII-BglII fragment containing the HBV pre-S1, pre-S2 and S coding regions was isolated from HBV-containing DNA. The 2.3 kb fragment was ligated together with the 4.5 kb fragment (obtained as described in 1) containing the methallothionein promoter.

2  $\mu$ l of the 2.3 kb fragment was mixed with 3  $\mu$ l of the 4.5 kb fragment and ligated together in a total volume of 10  $\mu$ l ligation buffer, containing 2 units T4-DNA ligase and 2 mM ATP at 14° C overnight.

The ligation mixture was added to 150  $\mu$ l competent bacterial cell suspension for DNA up-take. After the DNA up-take the bacterial cells were spread on LB agar plate containing 50  $\mu$ g/ml ampicillin at volumes of 50 to 300  $\mu$ l cell suspension per plate. The agar plates were incubated at 37° C overnight. Single isolated bacterial colonies were screened for the presence of a plasmid containing the desired fragment.

12. Conversion of a part of the HBV-gene sequence with HBV -core epitopes.

The plasmid resulting from 11 above was digested with the endonucleases BglII and XbaI. Two molecules were expected, one 550 bp fragment and a 6.25 kb fragment which was isolated after agarose gel electrophoresis.

The 6.25 kb fragment was ligated together with the 270 bp fragment (after digestion with BglII and XbaI and fragment isolation as described above) of the plasmid described in 10, coding for an epitope part of the HBV-core gene.

The ligation mixture was added to 150  $\mu$ l competent bacterial cell suspension for DNA up-take. Single isolated bacterial colonies were screened for the presence of the desired plasmid. The new plasmid was proved by a digestion with BamHI. Three molecules were expected, a 950 bp, a 450 bp and a 5,150 bp fragment.

13. Preparation of a "vehicle" plasmid.

The plasmid (described in 11) was digested with EcoRI and XbaI. Two molecules were expected, one 2,450 bp fragment and a 4,350 bp fragment which was isolated after gel electrophoresis.

This 4,350 bp fragment was ligated together with the oligo-DNA-nucleotide 39 (SEQ ID No.:29) coding for the entire DNA-sequence of HBV-S-gene from ATG to the XbaI site, wherein the ATG was changed into ATA.

14. Core-epitope upstream of the entire HBV-S gene.

This "vehicle" plasmid was then digested with PstI and XbaI, two molecules were expected, one 600 bp plasmid residue and a 3,850 bp fragment which was isolated and ligated together with a PstI-XbaI fragment of 2,800 bp (2,700 bp) isolated after digestion of the plasmid described in 10.

After screening for the desired plasmid, it was characterized by restriction endonuclease digestion with EcoRI and XbaI, EcoRI and BglII and BamHI.

15. Insertion of a selection marker.

The plasmid (described in 14) was linearized with EcoRI. The reaction was performed in a total volume of 50 µl and a final concentration of 1 µg/µl plasmid DNA. 50 units of EcoRI were added and the digestion was proved after incubation at 37° C for three hours by agarose gel electrophoresis.

The reaction was stopped by adding 1 µl of 0.5 M EDTA and DNA was precipitated with a final concentration of 0.3 M sodium acetate and 3-4 volumes of ethanol at -80° C for 30 minutes. The precipitated DNA was dissolved in 50 µl distilled water.

2 µl of the linearized plasmid was mixed with 3 µl of the DNA fragment containing the methallothionein promoter and the neomycin selection gene (described in 4) and ligated together. Single bacterial colonies were screened for the desired plasmid which was isolated, purified and characterized.

Each gene construct described above can be constructed also with the U2-promoter whereby the MT-promoter-containing DNA fragment, after digestion with EcoRI and BglII, is replaced by a DNA fragment containing the U2-promoter isolated after digestion with EcoRI and BglII.

16. Isolation of the E coli xanthine quanine phosphoribosyl transferase (egpt) selection gene.

The fragment containing the egpt selection gene was isolated after digestion of the plasmid pMSG with BamHI and BglII (1.8kb) and ligated together with a 4.5 kb fragment (BglII-BamHI, described in 1) containing the MT-promoter. After screening for the desired plasmid it was isolated, purified and finalized by a conversion of the BamHI site into an EcoRI site.

17. Isolation of desired DNA sequences by PCR-method.

One DNA fragment (400 bp) was isolated after gel electrophoresis. It was generated by PCR-method (described in Example 5) by using the specific oligonucleotides 131 and 132 (SEQ ID No.:33 and 34) as primers.

The DNA fragment was digested with the endonucleases BamHI and XbaI and then purified by gel electrophoresis. The isolated PCR-fragment was ligated together with a 6.25 kb fragment which was isolated from the plasmid (described in 13) after digestion with BglII and XbaI. After DNA up-take and bacterial transformation the single bacterial colonies were screened for the desired plasmid.

18. Insertion of a selection marker.

The plasmid (described in 17) was finalized by adding a selection gene to the plasmid (described in 15).

19. Isolation of the H2K promoter.

The H2K promoter was isolated as an EcoRI and BglII fragment (2kb) from pSP65H2 (available from Exogene).

In all constructs described all promoters are replaceable as EcoRI/BglII fragments.

20. Conversion of a part of the HBV-gene sequence.

The plasmid resulting from 11) above was digested with the endonucleases BglII and XbaI. Two molecules were expected, one of which is a 6.250 kb fragment which was isolated after agarose gel electrophoresis.

The 6.250 kb fragment was ligated together with oligo-DNA-nucleotide 23 (SEQ ID No.:28). The ligation mixture was added to 150 µl competent bacterial cell suspension for DNA up-take. Single isolated bacterial colonies were screened for the presence of the desired plasmid. The new plasmid was proven by a digestion with the endonucleases EcoRI and BglII. Two molecules were expected, one 1,9 kb and one 4.450 kb.

21. Insertion of a egpt selection marker.

The plasmid (described in 20) was linearized with EcoRI. The reaction was performed in a total volume of 100 µl and a final concentration of 0.6 µg/µl plasmid DNA. 60 units of EcoRI were added and the digestion was proved after incubation at 37° C for three hours by agarose gel electrophoresis. The reaction was stopped by adding 2 µl of 0.5 M EDTA and the DNA was precipitated with a final concentration of 0.3 M sodium acetate and 4 volumes of ethanol at -80° C for 1 hour. The precipitated DNA was dissolved in 50 µl distilled water.

2 µl of the linearized plasmid was mixed with 3 µl of the DNA-fragment (3.7 kb) containing the methallothionein promoter and the egpt selection gene (described in 16) by digestion with EcoRI and ligated together. Single colonies were screened for the presence of the desired plasmid. Each of the

described gene constructs in Table III are preparable in the same way as described above.

#### Example 4

##### 5 Transfection of Mammalian Cells with Constructs of the Present Invention.

In order to achieve secretion of substantial amounts of the HBV peptides encoded by constructs of the present invention, mammalian cells must be transfected with a DNA construct of the present invention. The cotransfection was performed in two steps (i.e. a separate transfection for each construct) or in a single step  
10 (i.e. one transfection using preparation of both constructs). Cotransfection was confirmed either by use of different selection markers on the two constructs or by detection of secretion of expression products of both constructs by immunoassay.

Alternatively, a sequence encoding the HBV peptide sequence of the present invention and a separate sequence encoding the entire S or core or HAV protein could be combined in a single construct.

#### Example 5

##### Polymerase chain reaction (PCR).

20 The polymerase chain reaction allows to amplify specific DNA nucleotide sequences of a selected region of a known genomic sequence in vitro by more than a millionfold (Thomas J. White, Norman Arnheim, Henry A. Erlich 1989: The polymerase chain reaction. Technical Focus, Vol. 5. No. 6; S. Kwok and R. Higuchi 1989: Avoiding false positives with PCR. Nature, Vol. 339, pp 237-238).

DNA isolated from cells or plasmid DNA is treated to separate its complementary strands. These  
25 strands are then annealed with an excess of two DNA oligonucleotides (each 20 - 25 base pairs long) that have been chemically synthesized to match sequences separated by X nucleotides (where X is generally between 50 to 2,000 base pairs).

The two oligonucleotides serve as specific primers for in vitro DNA synthesis catalysed by DNA polymerase which copies the DNA between the sequences corresponding to the two oligonucleotides. If the  
30 two primer oligonucleotides contain the correct sequence it is possible to create new digestion sites at the 5' and 3'.

After multiple cycles of reaction, a large amount of a DNA fragment of the desired length was obtained, purified by gel electrophoresis and characterized by restriction enzyme digestion and agarose gel elec-  
35 trophoresis. The amplified, purified DNA fragment was then used to ligate it together with other fragments i.e. plasmid.

The PCR-DNA fragments were amplified with blunt end. To get sticky end (for the ligation procedure) the fragment has to be digested with the desired endonucleases and purified again.

The PCR-reaction will work for 20 to 30 cycles. One cycle is separated into three steps with different reaction times and different reaction temperatures which is controlled by a PCR-thermo-cycler. The first  
40 step is "Denaturation" of the matrix-DNA (1 min/95° C), the second step is "Hybridisation" of matrix DNA and primers (1 min/55° C) followed by "Polymerisation" (2 min/72° C).

The final volume for one assay is 30 µl for example, which contains the following final concentrations: PCR-buffer (1 x), nucleotide-mix with 200 µM of each of the four nucleotides, 200ng for 30 µl of each of the two primers, 0.5 units Taq-Polymerase per 30 µl aqua bidest.

#### Example 6

##### Culturing of Transfected Cells to Secrete Protein

50 The recipient cells (C127 or CHO-cells available from ATCC) were seeded in normal growth medium (DMEM + 10% Fetal Calf Serum, Glucose and Glutamine) into petri-dishes (1-2 x 10<sup>6</sup> cells per dish, φ 10 cm) at day 1. The next day the medium was removed (4 hours before the DNA precipitate was added onto the cells), and the cells were washed twice with 1 x PBS. Then 8 ml DMEM without FCS were added, 4 hours later the DNA precipitate (prepared as described below) was added to the cells. Again after 4 hours  
55 the medium was removed, 3 ml of Glycerol-Mix (50 ml 2 x TBS buffer, 30 ml glycerol, 120 ml distilled water) were added. The Glycerol-Mix was immediately removed after an incubation at 37° C for 3 minutes and the cells were washed with 1 x PBS. The cells were cultivated overnight with 8 ml of DMEM with 10% FCS.

After 48 hours, the cells were recovered from the dish by treating with Trypsin-EDTA-Solution (0.025% Trypsin + 1mM EDTA). Afterwards, to remove the Trypsin-EDTA the cells were washed with 1 x PBS, suspended in DMEM with 10% FCS and distributed into 24 costar-well-plates (cells from one dish into four 24-well-plates).

When the cells had grown well, selection medium was added (concentration 0.5 - 1 mg/ml of neomycin or: xanthine (250 µg/ml), hypoxanthine (15 µg/ml) or adenine (25 µg/ml), thymidine (10 µg/ml), aminopterin (2 µg/ml), mycophenolic acid (25 µg/ml) for eco-gpt, for example). The medium was changed every week. The first growing cell colonies were seen after 2 weeks.

To 10 µg of plasmid DNA and 20 µg of carrier-DNA (salmon sperm DNA, calf-thymus DNA) TE-buffer (10 mM Tris-HCl, 1 mM EDTA, pH 7.05) was added to a final volume of 440 µl and mixed together with 60 µl 2 M CaCl<sub>2</sub>. Then the same amount of 2x TBS (Hepes 50 mM, NaCl 280 mM, Na<sub>2</sub>HPO<sub>4</sub> 1.5 mM, pH 7.05) was added and mixed well. The precipitation solution was incubated for 30 minutes at 37° C and added directly to the cells which were to be transfected.

#### Example 7

Preparation of the Adjuvant of Purified Particles.

To the desired concentration of antigen suspended in sterile saline, 1 : 10,000 volume Thimerosol, 1/10 volume of filter-sterilized 0.2 M KAl(SO<sub>4</sub>)<sub>2</sub> · 12 H<sub>2</sub>O were added. The pH was adjusted to 5.0 with sterile 1 N NaOH and the suspension was stirred at room temperature for 3 hours. The alum-precipitated antigen was recovered by centrifugation for 10 minutes at 2,000 rpm, resuspended in sterile normal saline containing 1:10,000 Thimerosol and aliquoted under sterile conditions.

#### Example 8

Purification of Hepatitis-B-core Antigen.

The cell supernatant of HB-core-antigen-secreting cells was collected and concentrated by ultrafiltration. The concentrate was cleared by centrifugation at 20,000 rpm for 15 minutes at 4° C in a Beckman SW28 rotor.

Particle formation was tested by sucrose density centrifugation (0-45% sucrose) in a Beckman SW28 rotor for 24 hours at 28,000 rpm and 4° C. The gradient was fractionated and the single fractions were analyzed by Elisa.

#### Example 9

The following tables give some results of Elisa analysis of immunogenic particles of the present invention as described below:

Table IV shows the Elisa data of the purified HBs-antigen particle produced from any HBV-sequence construct of the present invention including the pre-S1 epitopes and the S region with the anti-pre-S1 monoclonal antibody MA 18/7 and the anti-HBs monoclonal antibody G022.

Table IV shows the fractions (21) collected after CsCl density gradient.

Table IV-1

CsCl-gradient Fraction No.	Elisa Measurement (E = 492) Monoclonal Antibody 18/7
13	0.092
14	0.210
15	0.388
16	1.662
17	2.604
18	0.648
19	0.031

Table IV-2

CsCl-gradient Fraction No.	Elisa Measurement (E = 492) Monoclonal Antibody G022
13	0.136
14	0.426
15	0.822
16	1.970
17	2.954
18	0.967
19	0.076

Table V shows the Elisa data of the purified HB-core-antigen particles produced from any HB-core-sequence construct of the present invention with polyclonal antibodies against HB-core and with monoclonal antibody G022 against HB-S-Ag.

Table V-1

Sucrose Gradient Fraction No.	Elisa Measurement (E = 492) Polyclonal Antibodies
6	0.25
7	0.922
8	1.423
9	1.5
10	1.5
11	1.28
12	0.466

Table V-2

Sucrose Gradient Fraction No.	Elisa Measurement (E = 492) Monoclonal Antibody G022
6	0.020
7	0.024
8	0.018
9	0.011
10	0.015
11	0.020
12	0.022

#### Example 10

Studies of administering Hepa-Care in Chimpanzees:

Hepa-Care are particles presenting hepatitis B surface antigens (S1 and S) in a specific formulation (ratio 50:50), which are used for the treatment of chronic carriers of hepatitis virus.

#### Experiment 1

A Hepatitis-B-carrier chimpanzee 1 was treated (intramuscularly) with Hepa-Care at time 0, 4, and 8 weeks with a dosage of 18  $\mu$ g per injection.

The liver enzymes were monitored (Fig. IV) as well as the hepatitis-B antigen level (Fig. V).



## Experiment 2

Chimpanzee 1 after treatment described above was given a booster treatment at week 30, 34, and 38. The results are shown in Fig. VI.

## Experiment 3

Chimpanzee 2 was treated with Hepa-Care, but contrary to chimpanzee 1 it was given intravenously. The dosage was 2 mg. The results are shown in Fig. VII.

From a control chimpanzee 3 the liver enzymes were also monitored and shown in Fig. VIII.

## Example 11

Treatment with Hepa-Care:

(for definition see Example 10)

Patient 1 (male, age = 65 years, disease for 2 years): Hepatitis-B parameters:

HBsAg	pos.
anti-HBs	neg.
HBeAg	neg.
anti-HBe	pos.
anti-HBc	neg.

was treated (i.m.) with Hepa-Care at month 0, 1, 6, and 7. The results of the antigen and antibody measurements are given in Fig. IX and X.

Patient 2 (female, age = 48 years, disease for 12 years):

Hepatitis-B parameters:

HBsAg	pos.
HBeAg	neg.
anti-HBs	neg.
anti-HBe	pos.
anti-HBc	pos.

was treated (i.m.) with Hepa-Care at month 0, 1, and 6. Results of antigen and antibody measurements are shown in Fig. XI and XII.

Patient 3 (female, age = 41 years, disease for 5 years):

Hepatitis-B parameters:

HBsAg	pos.
HBeAg	neg.
anti-HBs	neg.
anti-HBe	pos.

was treated at month 0, 1, 2, and 5 with Hepa-Care (i.m.). The measured values of HBs antigen and anti-HBs antibodies are shown in Fig. XIII and XIV.

SEQUENCE LISTING

5

SEQ ID NO: 1  
 SEQ TYPE: Nucleotide  
 SEQUENCE LENGTH: 558 bp  
 STRANDEDNESS: single  
 TOPOLOGY: linear  
 MOLECULE TYPE: genomic DNA  
 ORIGINAL SOURCE: HBV core  
 IMMEDIATE EXPERIMENTAL  
 SOURCE: PCR-amplification

10

15

20

25

30

35

40

45

50

55

ATG	GAC	ATT	GAC	CCT	TAT	AAA	GAA	TTT	GGA	GCT	
ACT	GTG	GAG	TTA	CTC	TCG	TTT	TTG	CCT	TCT	GAC	
TTC	TTT	CCT	TCC	GTA	CGA	GAT	CTC	CTA	GAC	ACC	
GCC	TCA	GCT	CTG	TAT	CGA	GAA	GCC	TTA	GAG	TCT	
CCT	GAG	CAT	TGC	TCA	CCT	CAC	CAT	ACT	GCA	CTC	
AGG	CAA	GCC	ATT	CTC	TGC	TGG	GGG	GAA	TTG	ATG	
ACT	CTA	GCT	ACC	TGG	GTG	GGT	AAT	AAT	TTG	CAA	
GAT	CCA	GCA	TCC	AGA	GAT	CTA	GTA	GTC	AAT	TAT	
GTT	AAT	ACT	AAC	ATG	GGT	TTA	AAG	ATC	AGG	CAA	
CTA	TTG	TGG	TTT	CAT	ATA	TCT	TGC	CTT	ACT	TTT	
GGA	AGA	GAG	ACT	GTA	CTT	GAA	TAT	TTG	GTC	TCT	
TTC	GGA	GTG	TGG	ATT	CGC	ACT	CCT	CCA	GCC	TAT	
AGA	CCA	CCA	AAT	GCC	CCT	ATG	TTA	TCA	ACA	CTT	
CCG	GAA	ACT	ACT	GTT	GTT	AGA	CGA	CGG	GAC	CGA	
GGC	AGG	TCC	CCT	AGA	AGA	AGA	ACT	CCC	TCG	CCT	
CGC	AGA	CGT	AGA	TCT	CAA	TCG	CCG	CGT	CGC	AGA	
AGA	TCT	CAA	TCT	CGG	GAA	TCT	CAA	TGT	TAG		

SEQ ID NO: 2  
 SEQ TYPE: Nucleotide  
 SEQUENCE LENGTH: 504 bp  
 STRANDEDNESS: single  
 TOPOLOGY: linear  
 MOLECULE TYPE: genomic DNA  
 ORIGINAL SOURCE: HBV core  
 IMMEDIATE EXPERIMENTAL  
 SOURCE: PCR-amplification

```

    ATG  GAC  ATT  GAC  CCT  TAT  AAA  GAA  TTT  GGA  GCT
    ACT  GTG  GAG  TTA  CTC  TCG  TTT  TTG  CCT  TCT  GAC
    TTC  TTT  CCT  TCC  GTA  CGA  GAT  CTC  CTA  GAC  ACC
    GCC  TCA  GCT  CTG  TAT  CGA  GAA  GCC  TTA  GAG  TCT
    CCT  GAG  CAT  TGC  TCA  CCT  CAC  CAT  ACT  GCA  CTC
    AGG  CAA  GCC  ATT  CTC  TGC  TGG  GGG  GAA  TTG  ATG
    ACT  CTA  GCT  ACC  TGG  GTG  GGT  AAT  AAT  TTG  CAA
    GAT  CCA  GCA  TCC  AGA  GAT  CTA  GTA  GTC  AAT  TAT
    GTT  AAT  ACT  AAC  ATG  GGT  TTA  AAG  ATC  AGG  CAA
    CTA  TTG  TGG  TTT  CAT  ATA  TCT  TGC  CTT  ACT  TTT
    GGA  AGA  GAG  ACT  GTA  CTT  GAA  TAT  TTG  GTC  TCT
    TTC  GGA  GTG  TGG  ATT  CGC  ACT  CCT  CCA  GCC  TAT
    AGA  CCA  CCA  AAT  GCC  CCT  ATG  TTA  TCA  ACA  CTT
    CCG  GAA  ACT  ACT  GTT  GTT  AGA  CGA  CGG  GAC  CGA
    GGC  AGG  TCC  CCT  AGA  AGA  AGA  ACT  CCC  TCG  CCT
    CGC  AGA  CGT
  
```

EP 0 491 077 A1

SEQ ID NO: 3  
 SEQ TYPE: Nucleotide  
 SEQUENCE LENGTH: 504 bp  
 STRANDEDNESS: single  
 TOPOLOGY: linear  
 MOLECULE TYPE: genomic DNA  
 ORIGINAL SOURCE: HBV core  
 IMMEDIATE EXPERIMENTAL  
 SOURCE: PCR-amplification

20	ATG	GAC	ATT	GAC	CCT	TAT	AAA	GAA	TTT	GGA	GCT
	ACT	GTG	GAG	TTA	CTC	TCG	TTT	TTG	CCT	TCT	GAC
25	TTC	TTT	CCT	TCC	GTA	CGA	GAT	CTC	CTA	GAC	ACC
	GCC	TCA	GCT	CTG	TAT	CGA	GAA	GCC	TTA	GAG	TCT
	CCT	GAG	CAT	TGC	TCA	CCT	CAC	CAT	ACT	GCA	CTC
30	AGG	CAA	GCC	ATT	CTC	TGC	TGG	GGG	GAA	TTG	ATG
	ACT	CTA	GCT	ACC	TGG	GTG	GGT	AAT	AAT	TTG	CAA
35	GAT	CCA	GCA	TCC	AGA	GAT	CTA	GTA	GTC	AAT	TAT
	GTT	AAT	ACT	AAC	ATG	GGT	TTA	AAG	ATC	AGG	CAA
	CTA	TTG	TGG	TTT	CAT	ATA	TCT	TGC	CTT	ACT	TTT
40	GGA	AGA	GAG	ACT	GTA	CTT	GAA	TAT	TTG	GTC	TCT
	TTC	GGA	GTG	TGG	ATT	CGC	ACT	CCT	CCA	GCC	TAT
	AGA	CCA	CCA	AAT	GCC	CCT	ATG	TTA	TCA	ACA	CTT
45	CCG	GAA	ACT	ACT	GTT	GTT	AGA	CGA	CGG	GAC	CGA
	GGC	AGG	TCC	CCT	AGA	AGA	AGA	ACT	CCC	TCG	CCT
50	CGC	AGA	CGT								

EP 0 491 077 A1

SEQ ID NO: 4  
 SEQ TYPE: Nucleotide  
 SEQUENCE LENGTH: 534 bp  
 STRANDEDNESS: single  
 TOPOLOGY: linear  
 MOLECULE TYPE: genomic DNA  
 ORIGINAL SOURCE: HBV core  
 IMMEDIATE EXPERIMENTAL  
 SOURCE: PCR-amplification

```

      TCC AAC CTG TGC CTT GGG TGG CTT TGG GGC
    ATG GAC ATT GAC CCT TAT AAA GAA TTT GGA GCT
    ACT GTG GAG TTA CTC TCG TTT TTG CCT TCT GAC
    TTC TTT CCT TCC GTA CGA GAT CTC CTA GAC ACC
    GCC TCA GCT CTG TAT CGA GAA GCC TTA GAG TCT
    CCT GAG CAT TGC TCA CCT CAC CAT ACT GCA CTC
    AGG CAA GCC ATT CTC TGC TGG GGG GAA TTG ATG
    ACT CTA GCT ACC TGG GTG GGT AAT AAT TTG CAA
    GAT CCA GCA TCC AGA GAT CTA GTA GTC AAT TAT
    GTT AAT ACT AAC ATG GGT TTA AAG ATC AGG CAA
    CTA TTG TGG TTT CAT ATA TCT TGC CTT ACT TTT
    GGA AGA GAG ACT GTA CTT GAA TAT TTG GTC TCT
    TTC GGA GTG TGG ATT CGC ACT CCT CCA GCC TAT
    AGA CCA CCA AAT GCC CCT ATG TTA TCA ACA CTT
    CCG GAA ACT ACT GTT GTT AGA CGA CGG GAC CGA
    GGC AGG TCC CCT AGA AGA AGA ACT CCC TCG CCT
    CGC AGA CGT
  
```

EP 0 491 077 A1

SEQ ID NO: 5  
 SEQ TYPE: Nucleotide  
 SEQUENCE LENGTH: 534 bp  
 STRANDEDNESS: single  
 TOPOLOGY: linear  
 MOLECULE TYPE: genomic DNA  
 ORIGINAL SOURCE: HBV core  
 IMMEDIATE EXPERIMENTAL  
 SOURCE: PCR-amplification

```

20      TCC AAC CTG TGC CTT GGG TGG CTT TGG GGC
      ATG GAC ATT GAC CCT TAT AAA GAA TTT GGA GCT
      ACT GTG GAG TTA CTC TCG TTT TTG CCT TCT GAC
25      TTC TTT CCT TCC GTA CGA GAT CTC CTA GAC ACC
      GCC TCA GCT CTG TAT CGA GAA GCC TTA GAG TCT
      CCT GAG CAT TGC TCA CCT CAC CAT ACT GCA CTC
30      AGG CAA GCC ATT CTC TGC TGG GGG GAA TTG ATG
      ACT CTA GCT ACC TGG GTG GGT AAT AAT TTG CAA
      GAT CCA GCA TCC AGA GAT CTA GTA GTC AAT TAT
35      GTT AAT ACT AAC ATG GGT TTA AAG ATC AGG CAA
      CTA TTG TGG TTT CAT ATA TCT TGC CTT ACT TTT
40      GGA AGA GAG ACT GTA CTT GAA TAT TTG GTC TCT
      TTC GGA GTG TGG ATT CGC ACT CCT CCA GCC TAT
      AGA CCA CCA AAT GCC CCT ATG TTA TCA ACA CTT
45      CCG GAA ACT ACT GTT GTT AGA CGA CGG GAC CGA
      GGC AGG TCC CCT AGA AGA AGA ACT CCC TCG CCT
50      CGC AGA CGT
  
```

SEQ ID NO: 6  
 SEQ TYPE: Nucleotide  
 SEQUENCE LENGTH: 87 bp  
 STRANDEDNESS: single  
 TOPOLOGY: linear  
 MOLECULE TYPE: genomic DNA  
 ORIGINAL SOURCE: HBV core  
 IMMEDIATE EXPERIMENTAL  
 SOURCE: chemically synthesized

ATC CTC TGC TGG GGG GAA TGG ATG ACT CTA GCT  
 ACC TGG GTG GGC AAT AAT TTG GAA GAT CCA GCA  
 TCT AGG GAC CTT GTA GTA AAT

SEQ ID NO: 7  
 SEQ TYPE: Nucleotide  
 SEQUENCE LENGTH: 81 bp  
 STRANDEDNESS: single  
 TOPOLOGY: linear  
 MOLECULE TYPE: genomic DNA  
 ORIGINAL SOURCE: HBV core  
 IMMEDIATE EXPERIMENTAL  
 SOURCE: chemically synthesized

GAC ATT GAC CCT TAT AAA GAA TTT GGA GCT ACT  
 GTG GAG TTA CTC TCG TTT TTG CCT TCT GAC TTC  
 TTT CCT TCC GTC AGG

EP 0 491 077 A1

SEQ ID NO: 8  
 SEQ TYPE: Nucleotide  
 SEQUENCE LENGTH: 114 bp  
 STRANDEDNESS: single  
 TOPOLOGY: linear  
 MOLECULE TYPE: genomic DNA  
 ORIGINAL SOURCE: HBV core  
 IMMEDIATE EXPERIMENTAL  
 SOURCE: chemically synthesized

TCC AAC CTG TGC CTT GGG TGG CTT TGG GGC ATG  
 GAC ATT GAC CCT TAT AAA GAA TTT GGA GCT ACT  
 GTG GAG TTA CTC TCG TTT TTG CCT TCT GAC TTC  
 TTT CCT TCC GTC AGG

SEQ ID NO: 9  
 SEQ TYPE: Nucleotide  
 SEQUENCE LENGTH: 90 bp  
 STRANDEDNESS: single  
 TOPOLOGY: linear  
 MOLECULE TYPE: genomic DNA  
 ORIGINAL SOURCE: HBV core  
 IMMEDIATE EXPERIMENTAL  
 SOURCE: chemically synthesized

GAT CTC CTA GAC ACC GCC TCA GCT CTG TAT CGA  
 GAA GCC TTA GAG TCT CCT GAG CTA TGC TCA CCT  
 CAC CAT ACT GCA CTC AGG CAA GGT



EP 0 491 077 A1

SEQ ID NO: 10  
 SEQ TYPE: Nucleotide  
 SEQUENCE LENGTH: 261 bp  
 STRANDEDNESS: single  
 TOPOLOGY: linear  
 MOLECULE TYPE: genomic DNA  
 ORIGINAL SOURCE: HBV core  
 IMMEDIATE EXPERIMENTAL  
 SOURCE: chemically synthesized

ATG GAC ATT GAC CCT TAT AAA GAA TTT GGA GCT  
 ACT GTG GAG TTA CTC TCG TTT TTG CCT TCT GAC  
 TTC TTT CCT TCC GTC AGG GAT CTC CTA GAC ACC  
 GCC TCA GCT CTG TAT CGA GAA GCC TTA GAG TCT  
 CCT GAG CTA TGC TCA CCT CAC CAT ACT GCA CTC  
 AGG CAA GGT ATC CTC TGC TGG GGG GAA TGG ATG  
 ACT CTA GCT ACC TGG GTG GGC AAT AAT TTG GAA  
 GAT CCA GCA TCT AGG GAC CTT GTA GTA AAT

EP 0 491 077 A1

SEQ ID NO: 11  
 SEQ TYPE: Nucleotide  
 SEQUENCE LENGTH: 291 bp  
 STRANDEDNESS: single  
 TOPOLOGY: linear  
 MOLECULE TYPE: genomic DNA  
 ORIGINAL SOURCE: HBV core  
 IMMEDIATE EXPERIMENTAL  
 SOURCE: chemically synthesized

	TCC	AAC	CTG	TGC	CTT	GGG	TGG	CTT	TGG	GGC
	ATG	GAC	ATT	GAC	CCT	TAT	AAA	GAA	TTT	GGA
	ACT	GTG	GAG	TTA	CTC	TCG	TTT	TTG	CCT	TCT
	TTC	TTT	CCT	TCC	GTC	AGG	GAT	CTC	CTA	GAC
	GCC	TCA	GCT	CTG	TAT	CGA	GAA	GCC	TTA	GAG
	CCT	GAG	CTA	TGC	TCA	CCT	CAC	CAT	ACT	GCA
	AGG	CAA	GGT	ATC	CTC	TGC	TGG	GGG	GAA	TGG
	ACT	CTA	GCT	ACC	TGG	GTG	GGC	AAT	AAT	TTG
	GAT	CCA	GCA	TCT	AGG	GAC	CTT	GTA	GTA	AAT

EP 0 491 077 A1

SEQ ID NO: 12  
 SEQ TYPE: Nucleotide  
 SEQUENCE LENGTH: 123 bp  
 STRANDEDNESS: single  
 5 TOPOLOGY: linear  
 MOLECULE TYPE: genomic DNA  
 ORIGINAL SOURCE: HBV core  
 10 IMMEDIATE EXPERIMENTAL  
 SOURCE: chemically synthesized

15 ACC TGG GTG GGT AAT AAT TTG CAA GAT CCA GCA  
 TCC AGA GAT CTA GTA GTC AAT TAT GTT AAT ACT  
 20 AAC ATG GGT TTA AAG ATC AGG CAA CTA TTG TGG  
 TTT CAT ATA TCT TGC CTT ACT TTT

25  
 30 SEQ ID NO: 13  
 SEQ TYPE: Nucleotide  
 SEQUENCE LENGTH: 138 bp  
 35 STRANDEDNESS: single  
 TOPOLOGY: linear  
 MOLECULE TYPE: genomic DNA  
 40 ORIGINAL SOURCE: HBV core  
 IMMEDIATE EXPERIMENTAL  
 SOURCE: chemically synthesized

45  
 50 GCA TCC AGA GAT CTA GTA GTC AAT TAT GTT AAT  
 ACT AAC ATG GGT TTA AAG ATC AGG CAA CTA TTG  
 TGG TTT CAT ATA TCT TGC CTT ACT TTT GGA AGA  
 GAG ACT GTA CTT GAA TAT TTG GTC TCT TTC GGA  
 55 GTG TGG

EP 0 491 077 A1

SEQ ID NO: 14  
 SEQ TYPE: Nucleotide  
 SEQUENCE LENGTH: 822 bp  
 STRANDEDNESS: single  
 TOPOLOGY: linear  
 MOLECULE TYPE: genomic DNA  
 ORIGINAL SOURCE: HBV S2 ayw/adw  
 IMMEDIATE EXPERIMENTAL  
 SOURCE: HBV DNA

ATG	CAG	TGG	AAT	TCC	AGA	ACC	TTC	CAC	CAA	ACT	CTG
CAA	GAT	CCC	AGA	GTG	AGA	GGC	CTG	TAT	TTC	CCT	GCT
GGT	GGC	TCC	AGT	TCA	GGA	ACA	GTA	AAC	CCT	GTT	CTG
ACT	ACT	GCC	TCT	CCC	TTA	TCG	TCA	ATC	TTC	TCG	AGG
ATA	GAG	AAC	ATC	ACA	TCA	GGA	TTC	CTA	GGA	CCC	CTT
CTC	GTG	TTA	CAG	GCG	GGG	TTT	TTC	TTG	TTG	ACA	AGA
ATC	CTC	ACA	ATA	CCG	CAG	AGT	CTA	GAC	TCG	TGG	TGG
ACT	TCT	CTC	AAT	TTT	CTA	GGG	GGA	ACT	ACC	GTG	TGT
CTT	GGC	CAA	AAT	TCG	CAG	TCC	TCA	ACC	TCC	AAT	CAC
TCA	CCA	ACC	TCT	TGT	CCT	CCA	ACT	TGT	CCT	GGT	TAT
CGC	TGG	ATG	TGT	CTG	CGG	CGT	TTT	ATC	ATC	TTC	CTC
TTC	ATC	CTG	CTG	CTA	TGC	CTC	ATC	TTC	TTG	TTG	GTT
CTT	CTG	GAC	TAT	CAA	GGT	ATG	TTG	CCC	GTT	TGT	CCT
CTA	ATT	CCA	GGA	TCC	TCA	ACA	ACC	AGC	ACG	GGA	CCA
TGC	CGG	ACC	TGC	ATG	ACT	ACT	GCT	CAA	GGA	ACC	TCT
ATG	TAT	CCC	TCC	TGT	TGC	TGT	ACC	AAA	CCT	TCG	GAC
GGA	AAT	TGC	ACC	TGT	ATT	CCC	ATC	CCA	TCA	TCC	TGG
GCT	TTC	GGA	AAA	TTC	CTA	TGG	GAG	TGG	GCC	TCA	GCC
CGT	TTC	TCC	TGG	CTC	AGT	TTA	CTA	GTG	CCA	TTT	GTT
CAG	TGG	TTC	GTA	GGG	CTT	TCC	CCC	ACT	GTT	TGG	CTT
TCA	GTT	ATA	TGG	ATG	ATG	TGG	TAT	TGG	GGG	CCA	AGT
CTG	TAC	AGC	ATC	TTG	AGT	CCC	TTT	TTA	CCG	CTG	TTA
CCA	ATT	TTC	TTT	TGT	CTT	TGG	GTA	TAC	ATT		

EP 0 491 077 A1

SEQ ID NO: 15  
 SEQ TYPE: Nucleotide  
 5. SEQUENCE LENGTH: 99 bp  
 STRANDEDNESS: single  
 TOPOLOGY: linear  
 10 MOLECULE TYPE: genomic DNA  
 ORIGINAL SOURCE: HBV core  
 IMMEDIATE EXPERIMENTAL  
 15 SOURCE: chemically synthesized

20 TAT GTT AAT ACT AAC ATG GGT TTA AAG ATC AGG  
 CAA CTA TTG TGG TTT CAT ATA TCT TGC CTT ACT  
 TTT GGA AGA GAG ACT GTA CTT GAA TAT TTG GTC

25

30

35

40

45

50

55

EP 0 491 077 A1

SEQ ID NO: 16  
 SEQ TYPE: Nucleotide  
 SEQUENCE LENGTH: 390 bp  
 STRANDEDNESS: single  
 TOPOLOGY: linear  
 MOLECULE TYPE: genomic DNA  
 ORIGINAL SOURCE: HBV core  
 IMMEDIATE EXPERIMENTAL  
 SOURCE: PCR-amplification

	TCC	AAC	CTG	TGC	CTT	GGG	TGG	CTT	TGG	GGC
20	ATG	GAC	ATT	GAC	CCT	TAT	AAA	GAA	TTT	GGA
	ACT	GTG	GAG	TTA	CTC	TCG	TTT	TTG	CCT	TCT
	TTC	TTT	CCT	TCC	GTA	CGA	GAT	CTC	CTA	GAC
25	GCC	TCA	GCT	CTG	TAT	CGA	GAA	GCC	TTA	GAG
	CCT	GAG	CAT	TGC	TCA	CCT	CAC	CAT	ACT	GCA
	AGG	CAA	GCC	ATT	CTC	TGC	TGG	GGG	GAA	TTG
30	ACT	CTA	GCT	ACC	TGG	GTG	GGT	AAT	AAT	TTG
	GAT	CCA	GCA	TCC	AGA	GAT	CTA	GTA	GTC	AAT
35	GTT	AAT	ACT	AAC	ATG	GGT	TTA	AAG	ATC	AGG
	CTA	TTG	TGG	TTT	CAT	ATA	TCT	TGC	CTT	ACT
40	GGA	AGA	GAG	ACT	GTA	CTT	GAA	TAT	TTG	GTC

EP 0 491 077 A1

SEQ ID NO: 17  
SEQ TYPE: Nucleotide with corresponding protein  
5 SEQUENCE LENGTH: 60 bp  
STRANDEDNESS: single  
TOPOLOGY: linear  
10 MOLECULE TYPE: genomic DNA  
ORIGINAL SOURCE: HBV S1 ay  
IMMEDIATE EXPERIMENTAL  
SOURCE: chemically synthesized

15

20 AAT CCT CTG GGA TTC TTT CCC GAT CAC CAG TTG GAT

CCA GCC TTC AGA GCA AAC ACC GCA

25

30 Asn Pro Leu Gly Phe Phe Pro Asp His Gln Leu Asp

35 Pro Ala Phe Arg Ala Asn Thr Ala

40

45

50

55

EP 0 491 077 A1

SEQ ID NO: 18  
 SEQ TYPE: Nucleotide with corresponding protein  
 SEQUENCE LENGTH: 63 bp  
 STRANDEDNESS: single  
 TOPOLOGY: linear  
 MOLECULE TYPE: genomic DNA  
 ORIGINAL SOURCE: HBV S1 ay  
 IMMEDIATE EXPERIMENTAL  
 SOURCE: chemically synthesized

15

20

CCT GCC TCC ACC AAT CGC CAG TCA GGA AGG CAG - CCT

ACC CCG CTG TCT CCA CCT TTG AGA AAC

25

30

Pro Ala Ser Thr Asn Arg Gln Ser Gly Arg Gln Pro

Thr Pro Ile Ser Pro Pro Leu Arg Asn

35

40

45

50

55



SEQ ID NO: 19  
SEQ TYPE: Nucleotide with corresponding protein  
SEQUENCE LENGTH: 63 bp  
STRANDEDNESS: single  
TOPOLOGY: linear  
MOLECULE TYPE: genomic DNA  
ORIGINAL SOURCE: HBV S1 ay  
IMMEDIATE EXPERIMENTAL  
SOURCE: chemically synthesized

GAT CCA GCC TTC AGA GCA AAC ACC GCA AAT CCA GAT  
TGG GAC TTC AAT CCC AAC AAG GAC ACC

Asp Pro Ala Phe Arg Ala Asn Thr Ala Asn Pro Asp  
Trp Asp Phe Asn Pro Asn Lys Asp Thr

SEQ ID NO: 20  
SEQ TYPE: Nucleotide with corresponding protein  
SEQUENCE LENGTH: 108 bp  
5 STRANDEDNESS: single  
TOPOLOGY: linear  
MOLECULE TYPE: genomic DNA  
10 ORIGINAL SOURCE: HBV S1 ay  
IMMEDIATE EXPERIMENTAL  
SOURCE: chemically synthesized

15  
20 CCG CAC GGA GGC CTT TTG GGG TGG AGC CCT CAG GCT  
CAG GGC ATA CTA CAA ACT TTG CCA GCA AAT CCG CCT  
CCT GCC TCC ACC AAT CGC CAG TCA GGA AGG CAG CCT

25  
30 Pro His Gly Gly Leu Leu Gly Trp Ser Pro Gln  
Ala Gln Gly Ile Leu Glu Thr Leu Pro Ala Asn  
Pro Pro Pro Ala Ser Thr Asn Arg Gln Ser Gly  
35 Arg Gln Pro

EP 0 491 077 A1

SEQ ID NO: 21  
 SEQ TYPE: Nucleotide  
 SEQUENCE LENGTH: 63 bp  
 STRANDEDNESS: single  
 TOPOLOGY: linear  
 MOLECULE TYPE: genomic DNA  
 ORIGINAL SOURCE: HBV S1 ad  
 IMMEDIATE EXPERIMENTAL  
 SOURCE: chemically synthesized

15

CCT GCC TCC ACC AAT CGG CAG TCA GGA AGG CAG CCT

20

ACT CCC ATC TCT CCA CCT CTA AGA GAC - X

25

30

35

40

45

50

55

EP 0 491 077 A1

SEQ ID NO: 22  
 SEQ TYPE: Nucleotide with corresponding protein  
 SEQUENCE LENGTH: 108 bp  
 STRANDEDNESS: single  
 TOPOLOGY: linear  
 MOLECULE TYPE: genomic DNA  
 ORIGINAL SOURCE: HBV S1 ad  
 IMMEDIATE EXPERIMENTAL  
 SOURCE: chemically synthesized

CCA CAC GGC GGT ATT TTG GGG TGG AGC CCT CAG GCT  
 CAG GGC ATA TTG ACC ACA GTG TCA ACA ATT CCT CCT  
 CCT GCC TCC ACC AAT CGG CAG TCA GGA AGG CAG CCT  
 Pro His Gly Gly Ile Leu Gly Trp Ser Pro Gln  
 Ala Gln Gly Ile Leu Thr Thr Val Ser Thr Ile  
 Pro Pro Pro Ala Ser Thr Asn Arg Gln Ser Gly  
 Arg Gln Pro

EP 0 491 077 A1

SEQ ID NO: 23  
 SEQ TYPE: Nucleotide  
 SEQUENCE LENGTH: 60 bp  
 STRANDEDNESS: single  
 TOPOLOGY: linear  
 MOLECULE TYPE: genomic DNA  
 ORIGINAL SOURCE: HBV S2 ay  
 IMMEDIATE EXPERIMENTAL  
 SOURCE: chemically synthesized

CAG TGG AAT TCC AGA ACC TTC CAC CAA ACT CTG  
 CAA GAT CCC AGA GTG AGA GGC CTG TAT - X

SEQ ID NO: 24  
 SEQ TYPE: Nucleotide  
 SEQUENCE LENGTH: 60 bp  
 STRANDEDNESS: single  
 TOPOLOGY: linear  
 MOLECULE TYPE: genomic DNA  
 ORIGINAL SOURCE: HBV S2 ay  
 IMMEDIATE EXPERIMENTAL  
 SOURCE: chemically synthesized

GAT CCC AGA GTG AGA GGC CTG TAT TTC CCT GCT  
 GGT GGC TCC AGT TCA GGA ACA GTA AAC - X

EP 0 491 077 A1

SEQ ID NO: 25  
 SEQ TYPE: Nucleotide  
 SEQUENCE LENGTH: 558 bp  
 STRANDEDNESS: single  
 TOPOLOGY: linear  
 MOLECULE TYPE: genomic DNA  
 ORIGINAL SOURCE: HBV core adw  
 IMMEDIATE EXPERIMENTAL  
 SOURCE: PCR-amplification

	ATG	GAC	ATT	GAC	CCT	TAT	AAA	GAA	TTT	GGA	GCT
20	ACT	GTG	GAG	TTA	CTC	TCG	TTT	TTG	CCT	TCT	GAC
	TTC	TTT	CCT	TCC	GTA	CGA	GAT	CTC	CTA	GAC	ACC
	GCC	TCA	GCT	CTG	TAT	CGA	GAA	GCC	TTA	GAG	TCT
25	CCT	GAG	CAT	TGC	TCA	CCT	CAC	CAT	ACT	GCA	CTC
	AGG	CAA	GCC	ATT	CTC	TGC	TGG	GGG	GAA	TTG	ATG
30	ACT	CTA	GCT	ACC	TGG	GTG	GGT	AAT	AAT	TTG	CAA
	GAT	CCA	GCA	TCC	AGA	GAT	CTA	GTA	GTC	AAT	TAT
	GTT	AAT	ACT	AAC	ATG	GGT	TTA	AAG	ATC	AGG	CAA
35	CTA	TTG	TGG	TTT	CAT	ATA	TCT	TGC	CTT	ACT	TTT
	GGA	AGA	GAG	ACT	GTA	CTT	GAA	TAT	TTG	GTC	TCT
	TTC	GGA	GTG	TGG	ATT	CGC	ACT	CCT	CCA	GCC	TAT
40	AGA	CCA	CCA	AAT	GCC	CCT	ATG	TTA	TCA	ACA	CTT
	CCG	GAA	ACT	ACT	GTT	GTT	AGA	CGA	CGG	GAC	CGA
45	GGC	AGG	TCC	CCT	AGA	AGA	AGA	ACT	CCC	TCG	CCT
	CGC	AGA	CGT	AGA	TCT	CAA	TCG	CCG	CGT	CGC	AGA
	AGA	TCT	CAA	TCT	CGG	GAA	TCT	CAA	TGT	TAG	

EP 0 491 077 A1

SEQ ID NO: 26  
 SEQ TYPE: Nucleotide  
 SEQUENCE LENGTH: 678 bp  
 STRANDEDNESS: single  
 TOPOLOGY: linear  
 MOLECULE TYPE: genomic DNA  
 ORIGINAL SOURCE: HBV S adw/ayw  
 IMMEDIATE EXPERIMENTAL  
 SOURCE: HBV DNA

	ATA	GAG	AAC	ATC	ACA	TCA	GGA	TTC	CTA	GGA	CCC	CTT	CTC
20	GTG	TTA	CAG	GCG	GGG	TTT	TTC	TTG	TTG	ACA	AGA	ATC	CTC
	ACA	ATA	CCG	CAG	AGT	CTA	GAC	TCG	TGG	TGG	ACT	TCT	CTC
	AAT	TTT	CTA	GGG	GGA	ACT	ACC	GTG	TGT	CTT	GGC	CAA	AAT
25	TCG	CAG	TCC	TCA	ACC	TCC	AAT	CAC	TCA	CCA	ACC	TCT	TGT
	CCT	CCA	ACT	TGT	CCT	GGT	TAT	CGC	TGG	ATG	TGT	CTG	CGG
	CGT	TTT	ATC	ATC	TTC	CTC	TTC	ATC	CTG	CTG	CTA	TGC	CTC
30	ATC	TTC	TTG	TTG	GTT	CTT	CTG	GAC	TAT	CAA	GGT	ATG	TTG
	CCC	GTT	TGT	CCT	CTA	ATT	CCA	GGA	TCC	TCA	ACA	ACC	AGC
35	ACG	GGA	CCA	TGC	CGG	ACC	TGC	ATG	ACT	ACT	GCT	CAA	GGA
	ACC	TCT	ATG	TAT	CCC	TCC	TGT	TGC	TGT	ACC	AAA	CCT	TCG
	GAC	GGA	AAT	TGC	ACC	TGT	ATT	CCC	ATC	CCA	TCA	TCC	TGG
40	GCT	TTC	GGA	AAA	TTC	CTA	TGG	GAG	TGG	GCC	TCA	GCC	CGT
	TTC	TCC	TGG	CTC	AGT	TTA	CTA	GTG	CCA	TTT	GTT	CAG	TGG
	TTC	GTA	GGG	CTT	TCC	CCC	ACT	GTT	TGG	CTT	TCA	GTT	ATA
45	TGG	ATG	ATG	TGG	TAT	TGG	GGG	CCA	AGT	CTG	TAC	AGC	ATC
	TTG	AGT	CCC	TTT	TTA	CCG	CTG	TTA	CCA	ATT	TTC	TTT	TGT
	CTT	TGG	GTA	TAC	ATT								

EP 0 491 077 A1

SEQ ID NO: 27  
 SEQ TYPE: Nucleotide  
 SEQUENCE LENGTH: 585 bp  
 STRANDEDNESS: single  
 TOPOLOGY: linear  
 MOLECULE TYPE: genomic DNA  
 ORIGINAL SOURCE: HBV S adw/ayw  
 IMMEDIATE EXPERIMENTAL  
 SOURCE: HBV DNA

CTA GAC TCG TGG TGG ACT TCT CTC AAT TTT CTA GGG  
 GGA TCT CCC GTG TGT CTT GGC CAA AAT TCG CAG TCC  
 CCA ACC TCC AAT CAC TCA CCA ACC TCC TGT CCT CCA  
 ATT TGT CCT GGT TAT CGC TGG ATG TGT CTG CGG CGT  
 TTT ATC ATA TTC CTC TTC ATC CTG CTG CTA TGC CTC  
 ATC TTC TTA TTG GTT CTT CTG GAT TAT CAA GGT ATG  
 TTG CCC GTT TGT CCT CTA ATT CCA GGA TCA ACA ACA  
 ACC AGT ACG GGA CCA TGC AAA ACC TGC ACG ACT CCT  
 GCT CAA GGC AAC TCT ATG TTT CCC TCA TGT TGC TGT  
 ACA AAA CCT ACG GAT GGA AAT TGC ACC TGT ATT CCC  
 ATC CCA TCG TCC TGG GCT TTC GCA AAA TAC CTA TGG  
 GAG TGG GCC TCA GTC CGT TTC TCT TGG CTC AGT TTA  
 CTA GTG CCA TTT GTT CAG TGG TTC GTA GGG CTT TCC  
 CCC ACT GTT TGG CTT TCA GCT ATA TGG ATG ATG TGG  
 TAT TGG GGG CCA AGT CTG TAC AGC ATC GTG AGT CCC  
 TTT ATA CCG CTG TTA CCA ATT TTC TTT TGT CTC TGG  
 GTA TAC ATT



EP 0 491 077 A1

SEQ ID NO: 28  
 SEQ TYPE: Nucleotide  
 SEQUENCE LENGTH: 106 bp  
 STRANDEDNESS: single  
 TOPOLOGY: linear  
 MOLECULE TYPE: genomic DNA  
 ORIGINAL SOURCE: HBV S1  
 IMMEDIATE EXPERIMENTAL  
 SOURCE: chemically synthesized

3'-GAT-CTT-TAA-CAT-GGA-GAA-CAA-TCC-TCT-G  
 GG-ATT-CTT-TCC-CGA-TCA-CCA-GTT-GGA-TCC-A  
 GC-CTT-CAG-AGC-AAA-CAC-CGC-AAA-TCC-AGA-T  
 TG-GGA-CTT-CAA-TCC-CAG-(T)-3'

SEQ ID NO: 29  
 SEQ TYPE: Nucleotide  
 SEQUENCE LENGTH: 115 bp  
 STRANDEDNESS: single  
 TOPOLOGY: linear  
 MOLECULE TYPE: genomic DNA  
 ORIGINAL SOURCE: HBV S  
 IMMEDIATE EXPERIMENTAL  
 SOURCE: chemically synthesized

5'-AAT-TCT-AGA-CTC-GAG-TCT-GAA-CAT-AGA-G  
 AA-CAT-CAC-ATC-AGG-ATT-CCT-AGG-ACC-CCT-T  
 CT-CGT-GTT-ACA-GGC-GGG-GTT-TTT-CTT-GTT-G  
 AC-AAG-AAT-CCT-CAC-AAT,ACC-GCA-GAG-(C)-3'

EP 0 491 077 A1

SEQ ID NO: 30  
 SEQ TYPE: Nucleotide  
 SEQUENCE LENGTH: 108 bp  
 STRANDEDNESS: single  
 TOPOLOGY: linear  
 MOLECULE TYPE: genomic DNA  
 ORIGINAL SOURCE: HBV core  
 IMMEDIATE EXPERIMENTAL  
 SOURCE: chemically synthesized

5'-GAT-CTT-TTA-AAG-GGA-TCC-TCT-GCT-GGG-G  
 GG-AAT-GGA-TGA-CTC-TAG-CTA-CCT-GGG-TGG-G  
 CA-ATA-ATT-TGG-AAG-ATC-CAG-CAT-CTA-GGG-A  
 CC-TTG-TAG-TAA-ATC-TAG-AC-(A)-3'

SEQ ID NO: 31  
 SEQ TYPE: Nucleotide  
 SEQUENCE LENGTH: 106 bp  
 STRANDEDNESS: single  
 TOPOLOGY: linear  
 MOLECULE TYPE: genomic DNA  
 ORIGINAL SOURCE: HBV core  
 IMMEDIATE EXPERIMENTAL  
 SOURCE: chemically synthesized

5'-GAT-CTC-CGG-GAA-TTC-CTG-GGG-CAT-GGA-C  
 AT-TGA-CCC-TTA-TAA-GGA-ATT-TGG-AGC-TAC-T  
 GT-GGA-GTT-ACT-CTC-GTT-TTT-GCC-TTC-TGA-C  
 TT-CTT-TCC-TTC-CGT-CAG-(G)-3'

5      SEQ ID NO:                      32  
       SEQ TYPE:                     Nucleotide  
       SEQUENCE LENGTH:            89 bp  
       STRANDEDNESS:                single  
 10     TOPOLOGY:                     linear  
       MOLECULE TYPE:                genomic DNA  
       ORIGINAL SOURCE:             HBV core  
 15     IMMEDIATE EXPERIMENTAL  
       SOURCE:                        chemically synthesized

20  
       5'-GAT-CTC-CTA-GAC-ACC-GCC-TCA-GCT-CTG-T  
 25     AT-CGA-GAA-GCC-TTA-GAG-TCT-CCT-GAG-CAT-T  
       GC-TCA-CCT-CAC-CAT-ACT-GCA-CTC-AGG-CAA-G  
 30     -(G)-3'

35      SEQ ID NO:                      33  
       SEQ TYPE:                     Nucleotide  
       SEQUENCE LENGTH:            25 bp  
 40     STRANDEDNESS:                single  
       TOPOLOGY:                     linear  
       MOLECULE TYPE:                genomic DNA  
       ORIGINAL SOURCE:             HBV core  
 45     IMMEDIATE EXPERIMENTAL  
       SOURCE:                        chemically synthesized

50  
       5'-TTG-GAT-CCT-CCA-ACC-TGT-GCC-TTG-(G)-3'

EP 0 491 077 A1

SEQ ID NO: 34  
SEQ TYPE: Nucleotide  
5 SEQUENCE LENGTH: 25 bp  
STRANDEDNESS: single  
TOPOLOGY: linear  
10 MOLECULE TYPE: genomic DNA  
ORIGINAL SOURCE: HBV core  
IMMEDIATE EXPERIMENTAL  
15 SOURCE: chemically synthesized

20 5'-CCT-CTA-GAA-CCA-AAT-ATT-CAA-GTA-(C)-3'

25  
30  
35  
40  
45  
50  
55  
.

# EP 0 491 077 A1

SEQ ID NO: 35  
 SEQ TYPE: Nucleotide  
 SEQUENCE LENGTH: 588 bp  
 STRANDEDNESS: single  
 TOPOLOGY: linear  
 MOLECULE TYPE: genomic DNA  
 ORIGINAL SOURCE: HBV core  
 IMMEDIATE EXPERIMENTAL  
 SOURCE: PCR-amplification

```

      TCC AAC CTG TGC CTT GGG TGG CTT TGG GGC
20  ATG GAC ATT GAC CCT TAT AAA GAA TTT GGA GCT
    ACT GTG GAG TTA CTC TCG TTT TTG CCT TCT GAC
25  TTC TTT CCT TCC GTA CGA GAT CTC CTA GAC ACC
    GCC TCA GCT CTG TAT CGA GAA GCC TTA GAG TCT
    CCT GAG CAT TGC TCA CCT CAC CAT ACT GCA CTC
30  AGG CAA GCC ATT CTC TGC TGG GGG GAA TTG ATG
    ACT CTA GCT ACC TGG GTG GGT AAT AAT TTG CAA
    GAT CCA GCA TCC AGA GAT CTA GTA GTC AAT TAT
35  GTT AAT ACT AAC ATG GGT TTA AAG ATC AGG CAA
    CTA TTG TGG TTT CAT ATA TCT TGC CTT ACT TTT
    GGA AGA GAG ACT GTA CTT GAA TAT TTG GTC TCT
40  TTC GGA GTG TGG ATT CGC ACT CCT CCA GCC TAT
    AGA CCA CCA AAT GCC CCT ATG TTA TCA ACA CTT
45  CCG GAA ACT ACT GTT GTT AGA CGA CGG GAC CGA
    GGC AGG TCC CCT AGA AGA AGA ACT CCC TCG CCT
    CGC AGA CGT AGA TCT CAA TCG CCG CGT CGC AGA
50  AGA TCT CAA TCT CGG GAA TCT CAA TGT TAG
  
```

## Claims

1. A combination of
  - a) at least one polypeptide sequence mediating the antigenicity of one or more epitopes and
  - b) a carrier capable of presenting the epitope sequence(s) a), wherein the polypeptide sequence(s)

a) can be bound to carrier b) by adsorption, any chemical bonding or secondary valences.

2. A combination according to claim 1, characterised in that polypeptide sequence(s) a) mediate(s) the antigenicity of a T cell-activating epitope in direct or indirect way.
3. A combination according to claim 1 or 2, characterised in that said polypeptide sequence(s) a) is a polypeptide of hepatitis B virus.
4. A combination according to one of claims 1 to 3, characterised in that polypeptide sequence(s) a) is the amino acid sequence of one or more members selected from the group comprising the HB viral peptides pre S1, pre S2, S and the core antigens.
5. A combination according to one of claims 1 to 4, characterised in that the polypeptide sequence(s) a) may be modified:
  - i) by having arbitrary deletions, whereby an epitope comprising at least six consecutive amino acid residues is preserved,
  - ii) by having substitutions of one or several amino acids, or
  - iii) by carrying an additional amino acid sequence either at its N-terminus, at its C-terminus or as an insertion into the polypeptide sequence(s) a).
6. A combination according to one of claims 1 to 5, characterised in that polypeptide sequence(s) a) is myristylated.
7. A combination according to one of claims 1 to 6, characterised in that the carrier b) is a polysaccharide, a hydrophobic polymer or an inorganic molecule having particle form.
8. A combination according to one of claims 1 to 6, characterised in that the carrier b) is a second polypeptide sequence.
9. A combination according to claim 8, characterised in that polypeptide sequence b) upon secretion forms particles having a diameter of at least 10nm.
10. A combination according to claim 8 or 9, characterised in that polypeptide sequence b) is a substantial part of or the complete amino acid sequence of a polypeptide selected from a group comprising the HBV S-peptide, the HBV core-, the HAV core- and the HIV core-antigen as well as the surface antigens of poliovirus, HAV or HIV.
11. A combination according to one of claims 8 to 10, characterised in that polypeptide sequence b) may be modified:
  - i) by having arbitrary deletions, whereby the particle forming capacity is preserved,
  - ii) by having substitutions of one or several amino acids, or
  - iii) by carrying an additional amino acid sequence either at its N-terminus, at its C-terminus or as an insertion into the polypeptide sequence b).
12. A combination according to one of claims 8 to 11, characterised in that polypeptide sequence b) is myristylated.
13. A combination according to one of claims 1 to 6 or 8 to 12, characterised in that the polypeptide sequences a) and b) are linked via disulfide bridges.
14. A combination according to one of claims 1 to 6 or 8 to 13, characterised in that the polypeptide sequences a) and b) are linked via "hydrophobic anchoring" (mediated by myristic acid).
15. A combination according to one of claims 1 to 6 and 8 to 14, characterised in that the polypeptide sequences a) and b) are linked by a peptide bond, wherein optionally a spacer sequence is inserted in between polypeptide sequence(s) a) and polypeptide sequence b), said spacer sequence being linked via peptide bonds to polypeptide sequences a) and b).

16. Use of a combination according to any of the claims 1 to 15 for the production of a medicament for the treatment of chronic viral hepatitis.

17. A recombinant DNA molecule coding for a combination according to any of the claims 1 to 16 comprising at least one first DNA sequence, optionally a second, a third and/or a fourth DNA sequence, wherein

i) said at least one first DNA sequence codes for at least one polypeptide sequence a) according to one of claims 1 to 6.

ii) said second DNA sequence codes for a polypeptide sequence b) according to one of claims 8 to 12,

iii) said third DNA sequence codes for a spacer sequence according to claim 15, and

iv) said fourth DNA sequence codes for a selection marker,


and wherein the DNA sequences are controlled by DNA elements essential for expression, and optionally have a common reading frame.

18. A recombinant DNA molecule according to claim 17, characterised in that in the first DNA sequence(s) up to 30% nucleotides may be substituted.

19. A host cell transfected with a recombinant DNA molecule according to either of claims 17 or 18.


20. A host cell according to claim 19, characterised in that said host cell is selected from the group comprising mammalian, yeast and bacterial cells.

**Fig. 1** shows a DNA construct, coding for a promoter, a particle former sequence and a selection gene (described in Example 3/4)





**Fig. II** shows a DNA-gene construct containing a promoter, an epitope with the entire HB-S-Ag and a selection gene (described in Example 3/18)



**Fig. III** shows a DNA construct presenting a promoter, a T-cell epitope with a particle former residue and a selection gene (described in Example 3/21)






Fig. IV


**LIVER ENZYME MEASUREMENT**  
CHIMPANZEE EXPERIMENT 1



**FIG. V**      **ANTIGEN MEASUREMENT**  
**CHIMPANZEE EXPERIMENT 1**



**Fig. VI**      **LIVER ENZYME MEASUREMENT**  
**CHIMPANZEE EXPERIMENT 2**



**Fig. VII RESULTS OF CHIMPANZEE EXPERIMENT 3**




Fig. VIII Measurement of Liver Enzymes of a Control Chimpanzee without Treatment





Fig. IX


ANTIGEN MEASUREMENT

PATIENT #1







**Fig. X**                      **ANTIBODY MEASUREMENT**  
**PATIENT #1**




**Fig. XI**                      **ANTIGEN MEASUREMENT**  
**PATIENT #2**




**Fig. XII**                      **ANTIBODY MEASUREMENT**  
**PATIENT #2**



**Fig. XIII**                      **ANTIGEN MEASUREMENT**  
**PATIENT #3**



**Fig. XIV**                      **ANTIBODY MEASUREMENT**  
**PATIENT #3**





European Patent  
Office

## EUROPEAN SEARCH REPORT

Application Number

EP 90 12 4775

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.5)
X	EP-A-243 913 (CALIFORNIA INSTITUTE OF TECHNOLOGY) ) November 4, 1987 * the whole document *	1-5, 7-20	A61K39/12 A61K39/29 C12N15/33 C12N15/36
X	WO-A-8 810 300 (MEDICO LABS AG) * the whole document, especially examples 2 and 3 *	1-5, 8-20	
X	EP-A-271 302 (SCRIPPS CLINIC AND RESEARCH FOUNDATION) * the whole document *	1-5, 8-20	
X	EP-A-385 610 (THE WELLCOME FOUNDATION LIMITED) * the whole document *	1-5, 8-20	
X	EP-A-175 261 (CHIRON CORPORATION) * page 34 - page 39 *	1-3, 8-20	
X	EP-A-250 253 (SCRIPPS CLINIC AND RESEARCH FOUNDATION) * the whole document *	1-5, 8-20	
			TECHNICAL FIELDS SEARCHED (Int. Cl.5)
			C12N A61K C12P C07K
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 26 AUGUST 1991	Examiner CUPIDO, M
CATEGORY OF CITED DOCUMENTS			
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons A : member of the same patent family, corresponding document	

EPO FORM 1503 01.82 (P0001)